

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
7 November 2002 (07.11.2002)

PCT

(10) International Publication Number  
WO 02/088306 A2

- (51) International Patent Classification<sup>7</sup>: C12N (81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (21) International Application Number: PCT/US02/11853
- (22) International Filing Date: 26 April 2002 (26.04.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/287,539 30 April 2001 (30.04.2001) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): TSURUSHITA, Naoya [JP/US]; 3719 Redwood Circle, Palo Alto, CA 94306 (US). VASQUEZ, Maximiliano [CR/US]; 3818 Louis Road, Palo Alto, CA 94303 (US).
- (74) Agents: KELLEY, James, J. et al.; ELI LILLY AND COMPANY, P. O. Box 6288, Indianapolis, IN 46206-6288 (US).
- Published:  
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: HUMANIZED ANTIBODIES

(57) Abstract: Humanized forms of mouse antibody 3D6 that retain the binding properties of mouse 3D6 are disclosed. Also disclosed are processes for making the humanized antibody, intermediates for making the humanized antibodies, including, nucleotide sequences, vectors, transformed host cells, and methods of using the humanized antibody to treat, prevent, alleviate, reverse, or otherwise ameliorate symptoms or pathology or both, that are associated with Down's syndrome or pre-clinical or clinical Alzheimer's disease or cerebral amyloid angiopathy.

WO 02/088306 A2

## HUMANIZED ANTIBODIES

This application claims priority of US 60/287,539, filed 2001 April 30, the entire contents of which are incorporated herein by reference.

The invention relates to humanized antibodies useful for treating and preventing  
5 human diseases associated with amyloid  $\beta$  ( $A\beta$ ), such as Alzheimer's disease, Down's syndrome, and cerebral amyloid angiopathy. Mouse monoclonal antibody 3D6 has been widely used in analytical methods. After 3D6 was administered to a group of 11.5-12 month-old heterozygous, transgenic PDAPP mice ( $APP^{V717F}$ ) at a weekly intraperitoneal dose of about 10 mg/kg for six months, it has been reported that the mice had significantly  
10 reduced plaque burden, although the specific location of the reduction was not disclosed. [Bard, F., *et al.*, *Nature Med.* 6:916-919 (2000); WO 00/72876 and WO 00/72880, 7 December, 2000]. It was asserted that the antibody gained access to the central nervous system in sufficient amounts to "decorate"  $\beta$ -amyloid plaques. Finally, it was stated that mouse 3D6 induces phagocytosis of amyloid plaques in *in vitro* studies.

15 Methods for administering aggregated  $A\beta$ 1-42 to provoke an immunologic response and reduced amyloid deposits are described in PCT publication WO99/27944, published 10 June 1999. The description postulates that full-length aggregated  $A\beta$  peptide would be a useful immunogen. The application also indicates that antibodies that bind to  $A\beta$  peptide could be used as alternate therapeutic agents. However, this appears  
20 to be speculation since the supporting data reflect protocols that involve active immunization using, for example,  $A\beta$ 1-42.

WO 99/60024, published 25 November 1999, is directed to methods for amyloid removal using anti-amyloid antibodies. The mechanism, however, is stated to utilize the ability of anti- $A\beta$  antibodies to bind to pre-formed amyloid deposits (i.e. plaques) and  
25 result in subsequent microglial clearance of localized plaques. This mechanism was not proved *in vivo*. This publication further states that to be effective against  $A\beta$  plaques, anti- $A\beta$  antibodies must be delivered directly to the brain, because antibodies cannot cross the blood brain barrier.

Queen, *et al.* describe methods of humanizing antibodies [*e.g.*, US Patent Nos.  
30 5,585,089, 5,693,761, 5,693,762, 6,180,370].

Humanized forms of 3D6 are needed for use in humans having Down's syndrome, or pre-clinical or clinical Alzheimer's disease or cerebral amyloid angiopathy (CAA). However, it is not known whether 3D6 can be humanized so that the humanized antibody retained the binding properties of the mouse antibody.

5    Summary of the Invention

          This invention provides humanized forms of 3D6. These humanized antibodies have binding properties (affinity and epitope location) that are approximately the same as those of the mouse 3D6 antibody. The invention includes antibodies, single chain antibodies, and fragments thereof. The invention includes antibodies wherein the CDR  
10    are those of mouse monoclonal antibody 3D6 (sequences SEQ ID NO:1 through SEQ ID NO:6) and wherein the antibodies retain approximately the binding properties of the mouse antibody and have *in vitro* and *in vivo* properties functionally equivalent to the mouse antibody. In another aspect, this invention provides humanized antibodies and fragments thereof, wherein the variable regions have sequences comprising the CDR from  
15    mouse antibody 3D6 and specific human framework sequences (sequences SEQ ID NO:7 - SEQ ID NO:10), wherein the antibodies retain approximately the binding properties of the mouse antibody and have *in vitro* and *in vivo* properties functionally equivalent to the mouse antibody 3D6. In another aspect, this invention provides humanized antibodies and fragments thereof, wherein the light chain is SEQ ID NO:11 and the heavy chain is  
20    SEQ ID NO:12.

          Also part of the invention are polynucleotide sequences that encode the humanized antibodies or fragments thereof disclosed above, vectors comprising the polynucleotide sequences encoding the humanized antibodies or fragments thereof, host cells transformed with the vectors or incorporating the polynucleotides that express the humanized  
25    antibodies or fragments thereof, pharmaceutical formulations of the humanized antibodies and fragments thereof disclosed herein, and methods of making and using the same.

          Such humanized antibodies and fragments thereof are useful for, among other things, treating and preventing diseases and conditions characterized by A $\beta$  plaques or A $\beta$  toxicity in the brain, such as Alzheimer's disease, Down's syndrome, and cerebral  
30    amyloid angiopathy in humans.

The invention also includes use of a humanized antibody of the present invention for the manufacture of a medicament, including prolonged expression of recombinant sequences of the antibody or antibody fragment in human tissues, for treating, preventing, or reversing Alzheimer's disease, Down's syndrome, or cerebral amyloid angiopathy, or to inhibit the formation of amyloid plaques or the effects of toxic soluble A $\beta$  species in humans.

#### Detailed Description of the Invention

We have surprisingly found that humanized antibodies, wherein the CDRs originate from mouse monoclonal antibody 3D6 and the framework and other portions of the antibodies originate from a human germ line, bind A $\beta$ 1-40 and A $\beta$ 1-42 with at least the affinity with which mouse 3D6 binds A $\beta$ . Thus, we have a reasonable basis for believing that humanized antibodies of this specificity, modified to reduce their immunogenicity by converting them to a humanized form, offer the opportunity to treat, both prophylactically and therapeutically, conditions in humans that are associated with formation of beta-amyloid plaques. These conditions include, as noted above, pre-clinical and clinical Alzheimer's, Down's syndrome, and pre-clinical and clinical cerebral amyloid angiopathy.

As used herein, the word "treat" includes therapeutic treatment, where a condition to be treated is already known to be present and prophylaxis - *i.e.*, prevention of, or amelioration of, the possible future onset of a condition.

By "antibody" is meant a monoclonal antibody *per se*, or an immunologically effective fragment thereof, such as an Fab, Fab', or F(ab')<sub>2</sub> fragment thereof. In some contexts, herein, fragments will be mentioned specifically for emphasis; nevertheless, it will be understood that regardless of whether fragments are specified, the term "antibody" includes such fragments as well as single-chain forms. As long as the protein retains the ability specifically to bind its intended target, it is included within the term "antibody." Also included within the definition "antibody" are single chain forms. Preferably, but not necessarily, the antibodies useful in the invention are produced recombinantly. Antibodies may or may not be glycosylated, though glycosylated antibodies are preferred. Antibodies are properly cross-linked via disulfide bonds, as is well known.

The basic antibody structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function.

Light chains are classified as kappa and lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, and define the antibody's isotype as IgG, IgM, IgA, IgD and IgE, respectively. Within light and heavy chains, the variable and constant regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 3 or more amino acids.

The variable regions of each light/heavy chain pair form the antibody binding site. Thus, an intact antibody has two binding sites. The chains all exhibit the same general structure of relatively conserved framework regions (FR) joined by three hypervariable regions, also called complementarity determining regions or CDRs. The CDRs from the two chains of each pair are aligned by the framework regions, enabling binding to a specific epitope. From N-terminal to C-terminal, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with well known conventions [Kabat "Sequences of Proteins of Immunological Interest" National Institutes of Health, Bethesda, Md., 1987 and 1991; Chothia, et al., J. Mol. Biol. 196:901-917 (1987); Chothia, et al., Nature 342:878-883 (1989)].

By "humanized antibody" is meant an antibody that is composed partially or fully of amino acid sequences derived from a human antibody germline by altering the sequence of an antibody having non-human complementarity determining regions (CDR). A humanized immunoglobulin does not encompass a chimeric antibody, having a mouse variable region and a human constant region. However, the variable region of the antibody and even the CDR are humanized by techniques that are by now well known in the art. The framework regions of the variable regions are substituted by the corresponding human framework regions leaving the non-human CDR substantially intact. As mentioned above, it is sufficient for use in the methods of the invention, to

employ an immunologically specific fragment of the antibody, including fragments representing single chain forms.

Humanized antibodies have at least three potential advantages over non-human and chimeric antibodies for use in human therapy:

5           1) because the effector portion is human, it may interact better with the other parts of the human immune system (e.g., destroy the target cells more efficiently by complement-dependent cytotoxicity (CDC) or antibody-dependent cellular cytotoxicity (ADCC).

10           2) The human immune system should not recognize the framework or C region of the humanized antibody as foreign, and therefore the antibody response against such an injected antibody should be less than against a totally foreign non-human antibody or a partially foreign chimeric antibody.

15           3) Injected non-human antibodies have been reported to have a half-life in the human circulation much shorter than the half-life of human antibodies. Injected humanized antibodies will have a half-life essentially identical to naturally occurring human antibodies, allowing smaller and less frequent doses to be given.

The design of humanized immunoglobulins may be carried out as follows. As to the human framework region, a framework or variable region amino acid sequence of a CDR-providing non-human immunoglobulin is compared with corresponding sequences  
20           in a human immunoglobulin variable region sequence collection, and a sequence having a high percentage of identical amino acids is selected. When an amino acid falls under the following category, the framework amino acid of a human immunoglobulin to be used (acceptor immunoglobulin) is replaced by a framework amino acid from a CDR-providing non-human immunoglobulin (donor immunoglobulin):

25           (a) the amino acid in the human framework region of the acceptor immunoglobulin is unusual for human immunoglobulin at that position, whereas the corresponding amino acid in the donor immunoglobulin is typical for human immunoglobulin at that position;

          (b) the position of the amino acid is immediately adjacent to one of the CDRs; or

30           (c) any side chain atom of a framework amino acid is within about 5-6 angstroms (center-to-center) of any atom of a CDR amino acid in a three dimensional immunoglobulin model [Queen, *et al.*, Proc. Natl Acad. Sci. USA 86:10029-10033

(1989), and Co, *et al.*, Proc. Natl. Acad. Sci. USA 88, 2869 (1991)]. When each of the amino acid in the human framework region of the acceptor immunoglobulin and a corresponding amino acid in the donor immunoglobulin is unusual for human immunoglobulin at that position, such an amino acid is replaced by an amino acid typical for human immunoglobulin at that position.

A preferred humanized antibody is a humanized form of mouse antibody 3D6.

The CDRs of humanized 3D6 have the following amino acid sequences:

light chain CDR1:

1 5 10 15  
10 Lys Ser Ser Gln Ser Leu Leu Asp Ser Asp Gly Lys Thr Tyr Leu Asn  
(SEQ ID NO:1)

light chain CDR2:

1 5  
15 Leu Val Ser Lys Leu Asp Ser (SEQ ID NO:2)

light chain CDR3:

1 5  
20 Trp Gln Gly Thr His Phe Pro Arg Thr (SEQ ID NO:3)

heavy chain CDR1:

1 5  
Asn Tyr Gly Met Ser (SEQ ID NO:4)

25 heavy chain CDR2:

1 5 10 15  
Ser Ile Arg Ser Gly Gly Gly Arg Thr Tyr Tyr Ser Asp Asn Val Lys Gly  
(SEQ ID NO:5)

30 and, heavy chain CDR3:

1 5 10  
Tyr Asp His Tyr Ser Gly Ser Ser Asp Tyr (SEQ ID NO:6).

A preferred light chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline Vk segment DPK19 and J segment Jk4:

1 5 10 15  
40 Xaa Val Val Met Thr Gln Xaa Pro Leu Xaa Leu Pro Val Thr Xaa Gly  
20 25 30  
Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Asp Ser  
45 35 40 45  
Asp Gly Lys Thr Tyr Leu Asn Trp Leu Gln Gln Arg Pro Gly Gln Ser  
50 55 60

Pro Xaa Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro  
 65 70 75 80  
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
 5 85 90 95  
 Ser Arg Val Glu Ala Glu Asp Gly Val Tyr Tyr Cys Trp Gln Gly  
 100 105 110  
 10 Thr His Phe Pro Arg Thr Phe Gly Gly Thr Lys Xaa Glu Ile Lys  
 Arg (SEQ ID NO:7)

15

wherein:

Xaa at position 1 is Asp or Tyr;

Xaa at position 7 is Ser or Thr;

Xaa at position 10 is Ser or Thr;

20 Xaa at position 15 is Leu, Ile, or Val;

Xaa at position 50 is Arg or Lys;

Xaa at position 88 is Val or Leu; and

Xaa at position 109 is Val or Leu.

25 A preferred heavy chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline VH segment DP-45 and J segment JH4, with several amino acid substitutions to the consensus amino acids in the same human subgroup to reduce potential immunogenicity:

30 1 5 10 15  
 Glu Val Xaa Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 20 25 30  
 Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Asn Tyr  
 35 35 40 45  
 Gly Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 50 55 60  
 40 Ala Ser Ile Arg Ser Gly Gly Gly Arg Thr Tyr Tyr Ser Asp Asn Val  
 65 70 75 80  
 Lys Gly Arg Phe Thr Ile Ser Arg Glu Asn Ala Lys Asn Xaa Leu Tyr  
 45 85 90 95  
 Leu Gln Met Asn Ser Leu Xaa Xaa Glu Asp Thr Ala Val Tyr Tyr Cys



100                      105                      110  
 Val Arg Tyr Asp His Tyr Ser Gly Ser Ser Asp Tyr Trp Gly Gln Gly  
 115  
 5 Thr Xaa Val Thr Val Ser Ser  
 (SEQ ID NO:8)

wherein:

- Xaa at position 3 is Gln, Lys, or Arg;  
 10 Xaa at position 78 is Ser or Thr;  
 Xaa at position 87 is Arg or Lys;  
 Xaa at position 88 is Ala, Ser, or Thr; and  
 Xaa at position 114 is Leu, Thr, Ile, or Val.

- 15 A particularly preferred light chain variable region of a humanized antibody of the  
 present invention has the following amino acid sequence, in which the framework  
 originated from human germline Vk segment DPK19 and J segment Jk4:

1                      5                      10                      15  
 20 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly  
                     20                      25                      30  
                     Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Asp Ser  
 25                      35                      40                      45  
 Asp Gly Lys Thr Tyr Leu Asn Trp Leu Gln Gln Arg Pro Gly Gln Ser  
                     50                      55                      60  
 Pro Arg Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro  
 30                      65                      70                      75                      80  
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
                     85                      90                      95  
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Trp Gln Gly  
 35                      100                      105                      110  
 Thr His Phe Pro Arg Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys  
 40 Arg  
 (SEQ ID NO:9)

- A particularly preferred heavy chain variable region of a humanized antibody of  
 45 the present invention has the following amino acid sequence, in which the framework  
 originated from human germline VH segment DP-45 and J segment JH4: 1  
 5                      10                      15  
 Glu val Gln Leu val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

20 25 30  
 Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Asn Tyr  
 5 35 40 45  
 Gly Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 50 55 60  
 Ala Ser Ile Arg Ser Gly Gly Gly Arg Thr Tyr Tyr Ser Asp Asn Val  
 10 65 70 75 80  
 Lys Gly Arg Phe Thr Ile Ser Arg Glu Asn Ala Lys Asn Ser Leu Tyr  
 85 90 95  
 15 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 100 105 110  
 Val Arg Tyr Asp His Tyr Ser Gly Ser Ser Asp Tyr Trp Gly Gln Gly  
 115  
 20 Thr Leu Val Thr Val Ser Ser (SEQ ID NO:10).

A preferred light chain for a humanized antibody of the present invention has the  
 25 amino acid sequence:

1 5 10 15  
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly  
 20 25 30  
 30 Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Asp Ser  
 35 40 45  
 Asp Gly Lys Thr Tyr Leu Asn Trp Leu Gln Gln Arg Pro Gly Gln Ser  
 50 55 60  
 35 Pro Arg Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro  
 65 70 75 80  
 40 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
 85 90 95  
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Trp Gln Gly  
 100 105 110  
 45 Thr His Phe Pro Arg Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys  
 115 120 135  
 Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu  
 130 135 140  
 50 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe  
 145 150 155 160  
 Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln  
 165 170 175  
 55 Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser  
 180 185 190  
 60 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu

195 200 205  
Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser

5 210 215  
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys (SEQ ID NO:11).

A preferred heavy chain for a humanized antibody of the present invention has the  
10 amino acid sequence:

1 5 10 15  
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
20 25 30  
Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Asn Tyr  
35 40 45  
Gly Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
20 50 55 60  
Ala Ser Ile Arg Ser Gly Gly Gly Arg Thr Tyr Tyr Ser Asp Asn Val  
65 70 75 80  
Lys Gly Arg Phe Thr Ile Ser Arg Glu Asn Ala Lys Asn Ser Leu Tyr  
25 85 90 95  
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
100 105 110  
30 Val Arg Tyr Asp His Tyr Ser Gly Ser Ser Asp Tyr Trp Gly Gln Gly  
115 120 125  
Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe  
35 130 135 140  
Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu  
145 150 155 160  
40 Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp  
165 170 175  
Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu  
180 185 190  
45 Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser  
195 200 205  
Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro  
50 210 215 220  
Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys  
225 230 235 240  
55 Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro  
245 250 255  
Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser  
260 265 270  
60 Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp

```

      275              280              285
Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn

5      290              295              300
Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val

      305              310              315              320
Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu

10     325              330              335
Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys

      340              345              350
15    Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr

      355              360              365
Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr

20     370              375              380
Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu

      385              390              395              400
25    Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu

      405              410              415
Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys

      420              425              430
30    Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu

      435              440              445
Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly

35    Lys
                                           (SEQ ID NO:12)

```

Other sequences are possible for the light and heavy chains for humanized 3D6. The immunoglobulins can have two pairs of light chain/heavy chain complexes, at least one chain comprising one or more mouse complementarity determining regions

40 functionally joined to human framework region segments.

In another aspect, the present invention is directed to recombinant polynucleotides encoding antibodies which, when expressed, comprise the heavy and light chain CDRs from an antibody of the present invention. Exemplary polynucleotides, which on expression code for the polypeptide chains comprising the heavy and light chain CDRs of

45 monoclonal antibody 3D6 are given herein. Due to codon degeneracy, other polynucleotide sequences can be readily substituted for those sequences. Particularly preferred polynucleotides of the present invention encode antibodies, which when expressed, comprise the CDRs of SEQ ID NO:1 – SEQ ID NO:6, or any of the variable regions of SEQ ID NO:7 – SEQ ID NO:10, or the light and heavy chains of SEQ ID

50 NO:11 and SEQ ID NO:12.

The polynucleotides will typically further include an expression control polynucleotide sequence operably linked to the humanized immunoglobulin coding sequences, including naturally-associated or heterologous promoter regions. Preferably, the expression control sequences will be eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells, but control sequences for prokaryotic hosts may also be used. Once the vector has been incorporated into the appropriate host cell line, the host cell is propagated under conditions suitable for expressing the nucleotide sequences, and, as desired, the collection and purification of the light chains, heavy chains, light/heavy chain dimers or intact antibodies, binding fragments or other immunoglobulin forms may follow.

The nucleic acid sequences of the present invention capable of ultimately expressing the desired humanized antibodies can be formed from a variety of different polynucleotides (genomic or cDNA, RNA, synthetic oligonucleotides, etc.) and components (e.g., V, J, D, and C regions), using any of a variety of well known techniques. Joining appropriate genomic and synthetic sequences is a common method of production, but cDNA sequences may also be utilized.

Below is a cDNA sequence (SEQ ID NO:17), from which the light chain having the amino acid sequence of SEQ ID NO:19 may be expressed.

```

20      1  ATGATGAGTCCTGCCAGTTCTGTTTCTGTTAGTGCTCTGGATTCCGGAAACCAACGGT      60
      M M S P A Q F L F L L V L W I R E T N G

      61  GATGTTGTGATGACCCAGTCTCCACTCTCCTTGCCTGTTACCCTGGGACAACCAGCCTCC      120
25      D V V M T Q S P L S L P V T L G Q P A S

      121  ATCTCTTGCAAGTCAAGTCAGAGCCTCTTAGATAGTGATGGAAAGACATATTTGAATTGG      180
30      I S C K S S Q S L L D S D G K T Y L N W

      181  TTGCAACAGCGCCAGGCCAGTCTCCAAGACGCCTAATCTATCTGGTGTCTAAACTGGAC      240
      L Q Q R P G Q S P R R L I Y L V S K L D

      241  TCTGGAGTCCCTGACAGGTTCTCTGGCAGTGGATCAGGGACAGATTTTACACTGAAAATC      300
35      S G V P D R F S G S G S G T D F T L K I

      301  AGCAGAGTCGAGGCTGAGGATGTGGGAGTTTATTATTGCTGGCAAGGTACACATTTTCCT      360
40      S R V E A E D V G V Y Y C W Q G T H F P

      361  CGGACGTTGGTGGAGGCACCAAGGTGGAATCAAACGTAAGTGTGGCTGCACCATCTGTG      420
45      R T F G G G T K V E I K R T V A A P S V

      421  TTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCTG      480

```

```

      F I F P P S D E Q L K S G T A S V V C L
      CTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGAAGGTGGATAACGCCCTCCAA
5  481  -----+-----+-----+-----+-----+ 540
      L N N F Y P R E A K V Q W K V D N A L Q

      TCGGGTAACCTCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTC
10 541  -----+-----+-----+-----+-----+ 600
      S G N S Q E S V T E Q D S K D S T Y S L

      AGCAGCACCTTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAA
601  -----+-----+-----+-----+-----+ 660
      S S T L T L S K A D Y E K H K V Y A C E

      GTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT (SEQ ID NO:17)
15 661  -----+-----+-----+-----+-----+ 720
      V T H Q G L S S P V T K S F N R G E C (SEQ ID NO:19)

```

Below is a cDNA sequence (SEQ ID NO:18), from which the heavy chain having  
 20 the amino acid sequence of SEQ ID NO:20 may be expressed.

```

      ATGAACCTTCGGGCTCAGCTTGATTTTCCTTGTCTTCTTAAAGGTGTCCAGTGTGAA
1  1  -----+-----+-----+-----+-----+ 60
      M N F G L S L I F L V L V L K G V Q C E

      GTGCAACTGGTGGAGTCTGGGGGAGGCTTAGTGCAGCCTGGAGGCTCTCTGAGGCTCTCC
25 61  -----+-----+-----+-----+-----+ 120
      V Q L V E S G G G L V Q P G G S L R L S

      TGTGCAGGCTCTGGATTCACTTTTCAGTAACTATGGCATGTCTTGGGTTCCGCCAGGCTCCT
30 121 -----+-----+-----+-----+-----+ 180
      C A G S G F T F S N Y G M S W V R Q A P

      GGAAAGGGACTGGAGTGGGTTGCATCCATTAGGAGTGGTGGTGGTAGAACCTACTATTCA
35 181 -----+-----+-----+-----+-----+ 240
      G K G L E W V A S I R S G G G R T Y Y S

      GACAATGTAAAGGGCCGATTACCATCTCCAGAGAGAATGCCAAGAACAGCCTGTACCTG
40 241 -----+-----+-----+-----+-----+ 300
      D N V K G R F T I S R E N A K N S L Y L

      CAAATGAACAGTCTGAGAGCTGAGGACACGGCTGTCTATTATTGTGTGAGATATGATCAC
45 301 -----+-----+-----+-----+-----+ 360
      Q M N S L R A E D T A V Y Y C V R Y D H

      TATAGTGGTAGCTCCGACTACTGGGGCCAGGGCACCTTGGTCACAGTCTCCTCAGCCTCC
50 361 -----+-----+-----+-----+-----+ 420
      Y S G S S D Y W G Q G T L V T V S S A S

      ACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACA
55 421 -----+-----+-----+-----+-----+ 480
      T K G P S V F P L A P S S K S T S G G T

      GCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGTTGAAC
60 481 -----+-----+-----+-----+-----+ 540
      A A L G C L V K D Y F P E P V T V S W N

      TCAGGCGCCCTGACCAGCGGCGTGACACCTTCCCGGCTGTCTTACAGTCTCCTCAGGACTC
541 -----+-----+-----+-----+-----+ 600
      S G A L T S G V H T F P A V L Q S S G L

      TACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATC
601 -----+-----+-----+-----+-----+ 660
      Y S L S S V V T V P S S S L G T Q T Y I

```

```

      TGC AACGTGAATCACAAGCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCT
661 -----+-----+-----+-----+-----+-----+-----+-----+ 720
      C N V N H K P S N T K V D K K V E P K S
5
      TGTGACAAAACCTCACACATGCCACCGTGCCAGCACCTGAACTCCTGGGGGGACCGTCA
721 -----+-----+-----+-----+-----+-----+-----+-----+ 780
      C D K T H T C P P C P A P E L L G G P S
10
      GTCTTCCTCTTCCCCCAAACCCAGGACACCCCTCATGATCTCCCGACCCCTGAGGTC
781 -----+-----+-----+-----+-----+-----+-----+-----+ 840
      V F L F P P K P K D T L M I S R T P E V
15
      ACATGCGTGGTGGTGGACGTGAGCCAGAACCCCTGAGGTCAAGTTCAACTGGTACGTG
841 -----+-----+-----+-----+-----+-----+-----+-----+ 900
      T C V V V D V S H E D P E V K F N W Y V
20
      GACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCAGC
901 -----+-----+-----+-----+-----+-----+-----+-----+ 960
      D G V E V H N A K T K P R E E Q Y N S T
25
      TACCGTGTGGTCAGCGTCCTCACCGTCTGCACCAGGACTGGCTGAATGGCAAGGAGTAC
961 -----+-----+-----+-----+-----+-----+-----+-----+ 1020
      Y R V V S V L T V L H Q D W L N G K E Y
30
      AAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCATCGAGAAAACCATCTCCAAGGCC
1021 -----+-----+-----+-----+-----+-----+-----+-----+ 1080
      K C K V S N K A L P A P I E K T I S K A
35
      AAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGACC
1081 -----+-----+-----+-----+-----+-----+-----+-----+ 1140
      K G Q P R E P Q V Y T L P P S R D E L T
40
      AAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTG
1141 -----+-----+-----+-----+-----+-----+-----+-----+ 1200
      K N Q V S L T C L V K G F Y P S D I A V
45
      GAGTGGGAGAGCAATGGGCAGCCGAGAACAACTACAAGACCACGCTCCCGTGTGGAC
1201 -----+-----+-----+-----+-----+-----+-----+-----+ 1260
      E W E S N G Q P E N N Y K T T P P V L D
50
      TCCGACGGCTCCTTCTCCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAG
1261 -----+-----+-----+-----+-----+-----+-----+-----+ 1320
      S D G S F F L Y S K L T V D K S R W Q Q
55
      GGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAG
1321 -----+-----+-----+-----+-----+-----+-----+-----+ 1380
      G N V F S C S V M H E A L H N H Y T Q K
60
      AGCCTCTCCCTGTCTCCGGGTAAA (SEQ ID NO:18)
1381 -----+-----+-----+-----+-----+-----+-----+-----+ 1416
      S L S L S P G K (SEQ ID NO:20)

```

The complete sequence of a humanized 3D6 light chain gene with introns (located between MluI and BamHI sites, as in pVk-Hu3D6) is shown below (SEQ ID NO:15).

The nucleotide number indicates its position in pVk-Hu3D6. The Vk and Ck exons are translated in single letter code; the dot indicates the translation termination codon. The mature light chain starts at the double-underlined aspartic acid (D). The intron sequence is in italics. The polyA signal is underlined. The expressed light chain corresponds to

SEQ ID NO:11 when mature.

619 ACGCGTCCACCATGATGAGTCTCTGCCAGTTCCTGTTTCTGTTAGTGCTCTGGATTTCGGGAAACCAACGGTGATGTTGTG  
                   M M S P A Q F L F L L V L W I R E T N G D = V V  
 5 699 ATGACCCAGTCTCCACTCTCCTTGCTGTACCCTGGGACAACAGCCTCCATCTCTTGCAAGTCAAGTCAGAGCCTCTT  
                   M T Q S P L S L P V T L G Q P A S I S C K S S Q S L L  
 779 AGATAGTGATGGAAAGACATATTTGAATTGGTTGCAACAGCGCCAGGCCAGTCTCCAAGACGCCAATCTATCTGGTGT  
                   D S D G K T Y L N W L Q Q R P G Q S P R R L I Y L V  
 859 CTAAACTGGACTCTGGAGTCCCTGACAGGTTCTCTGGCAGTGGATCAGGGACAGATTTTACACTGAAAATCAGCAGAGTC  
                   S K L D S G V P D R F S G S G S G T D F T L K I S R V  
 10 939 GAGGCTGAGGATGTGGGAGTTTATTATTGTGCGCAAGGTACACATTTTCTCGGACGTTCCGGTGGAGGCACCAAGGTGGA  
                   E A E D V G V Y Y C W Q G T H F P R T F G G G T K V E  
 1019 AATCAACGTAAGTGCACTTTCCTTCTAGAAATTCTAAACTCTGAGGGGGTTCGGATGACGTGGCCATTCTTTGCCTAAAG  
                   I K R  
 1099 CATTGAGTTTACTGCAAGGTGCAAAAAGCATGCAAGCCCTCAGAATGGCTGCAAAAGAGCTCCAACAAAACAATTTAGAA  
 15 1179 CTTTATTAAGGAATAGGGGAAGCTAGGAAGAACTCAAAACATCAAGATTTTAAATACGCTTCTTGGTCTCCTTGCTAT  
 1259 AATTATCTGGGATAAGCATGCTGTTTCTGCTGTGCTTAACATGCCCTGTGATATCCGCAAACAACACACCCAAGGGC  
 1339 AGAAGTTTGTACTTAAACACCATCTGTTTGTGCTTCTTCTCAGGAACCTGTGGCTGCACCATCTGTCTTCATCTTCCCG  
   T V A A P S V F I F P  
 1419 CCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCTGCTGAATAACTTCTATCCAGAGAGGCCAAAGT  
 20 1499 ACAGTGGAGGTTGGATAACGCCCTCCAATCGGGTAACCTCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCT  
                   Q W K V D N A L Q S G N S Q E S V T E Q D S T  
 1579 ACAGCCTCAGCAGCACCTGACGCTGAGCAAGCAGACTACGAGAAACACAAAGTCTACGCTGCGAAGTCAACCATCAG  
                   Y S L S S T L T L S K A D Y E K H K V Y A C E V T H Q  
 25 1659 GGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTAGAGGGAGAAGTGCCCCACCTGCTCCTCAGTTC  
                   G L S S P V T K S F N R G E C •  
 1739 CAGCCTGACCCCTCCCATCCTTTGGCCTCTGACCCCTTTTCCACAGGGGACCTACCCCTATTGCGGTCTCCAGCTCAT  
 1819 CTTTACCTCACCCCTCTCCTCTCTTGGCTTAAATTATGCTAATGTTGGAGGAGAATGAATAAATAAA GTGAATCTTT  
 1899 GCACCTGTGGTTTCTCTCTTCTCCTCATTTAATAATTATTATCTGTTGTTTACCAACTACTCAATTTCTCTTATAAGGGA  
 30 1979 CTAATATGTAGTCATCCTAAGGCGCATACCATTTATAAAATCATCCTTCATTCTATTTTACCCATCATCCTCTGCA  
 2059 AGACAGTCTCCCTCAAACCCACAAGCCTTCTGTCTCACAGTCCCCTGGGCCATGGTAGGAGAGACTTGCTTCTTGT  
 2139 TCCCCCTCTCAGCAAGCCCTCATAGTCTTTTAAAGGGTGACAGGTCTTACAGTCATATATCCTTTGATTCAATTCCT  
 2219 GAGAATCAACCAAGCAAAATTTTCAAAGAGAAACCTGCTATAAAGAGAATCATTCATTGCAACATGATATAAAATAA  
 2299 CAACACAATAAAGCAATTAATAAACAACATAGGGAATGTTTAAAGTTCATCATGGTACTTAGACTTAATGGAATG  
 35 2379 CATGCCCTTATTATCTTTTAAACAGGTACTGAGGGACTCCTGTCTGCCAAGGGCCGTATTGAGTACTTTCCACAACCTA  
 2459 ATTTAATCCACACTATACTGTGAGATTAAAAACATTCATTAAATGTTGCAAGGTTCTATAAAGCTGAGAGACAATAT  
 2539 ATCTATAACTCAGCAATCCCACTTCTAGGATC (SEQ ID NO:15)

The complete sequence of a humanized 3D6 heavy chain gene with introns  
 40 (located between MluI and BamHI sites, as in pVg1-Hu3D6) is shown below (SEQ ID  
 NO:16). The nucleotide number indicates its position in pVg1-Hu3D6. The V<sub>H</sub> and C<sub>H</sub>  
 exons are translated in single letter code; the dot indicates the translation termination  
 codon. The mature heavy chain starts at the double-underlined glutamine (Q). The intron  
 sequences are in *italic*. The polyA signal is underlined. The expressed heavy chain  
 45 corresponds to SEQ ID NO:12 when mature.

619 ACGCGTCCACCATGAACTTCGGGCTCAGCTTGATTTTCCTTGTCTTAAAGGTGTCCAGTGTGAAGTGCAACTG  
                   M N F G L S L I F L V L V L K G V Q C E = V Q L  
 50 699 GTGGAGTCTGGGGGAGGCTTAGTGACGCTTGAGGCTCTCTGAGGCTCTCCTGTGAGGCTCTGGATTCACTTTTCAGTAA  
                   V E S G G G L V Q P G G S S L R L S C A G S G T F T S N  
 779 CTATGGCATGTCTTGGGTTGCCAGGCTCTGGAAGGAGTGGAGTGGGTTGCATCCATTAGGAGTGGTGGTGGTAGAA  
                   Y G M S W V R Q A P G K G L E W V A S I R S G G G R  
 859 CCTACTATTAGACAATGTAAGGGCCGATCACCATCTCCAGAGAGAATGCCAAGAACAGCCTGTACCTGCAAATGAAC  
                   T Y Y S D N V K G R F T I S R E N A K N S L Y L Q M N  
 55 939 AGTCTGAGAGCTGAGGACACGGCTGTCTATTATTGTGTGAGATATGATCACTATACTGGTAGCTCCGACTACTGGGGCCA  
                   S L R A E D T A V Y Y C V R Y D H Y S G S S D Y W G Q  
 1019 GGGCACCTTGGTCACAGTCTCCTCAGGTGAGTCTCACAACCTCTAGAGCTTTCTGGGGCAGGCCAGGCCTGACCTTGGC  
                   G T L V T V S S



1099 TTTGGGGCAGGGAGGGGGCTAAGGTGAGGCAGGTGGCGCCAGCCAGGTGCACACCCAATGCCCATGAGCCGAGACTGG  
 1179 ACGCTGAACCTCGCGGACAGTTAAGAAGCCAGGGGCTCTGCGCCCTGGGCCCAGCTCTGTCCACACCGCGGTACATG  
 1259 GCACCACCTCTCTTGCAGCCTCCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGC  
 A S T K G P S V F P L A P S S K S T S G G  
 5 1339 ACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCGAACCGGTGACGGTGTCTGGAACTCAGGCGCCCTGACCAG  
 T A A L G C L V K D Y F P E P V T V S W N S G A L T S  
 1419 CGGCGTGACACCTTCCCGGTGTCTACAGTCTCCTAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCA  
 G V H T F P A V L Q S S G L Y S L S S V V T V P S S  
 1499 GCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAAGCCAGCAACACCAAGGTGGACAAGAAAGTGGTGAGAGG  
 10 S L G T Q T Y I C N V N H K P S N T K V D K K V  
 1579 CCAGCACAGGGAGGGAGGGTGTCTGCTGGAAGCCAGGCTCAGCGCTCCTGCCTGGACGCATCCCGGCTATGCAGCCCCAG  
 1659 TCCAGGGCAGCAAGGCAGGCCCCGTCTGCCTCTTACCCGGAGGCTCTGCCCCCCCCACTCATGCTCAGGGAGAGGGTTC  
 1739 TTCTGGCTTTTTTCCCCAGGCTCTGGGCAGGCACAGGCTAGGTGCCCTTAACCCAGGCCCTGCACACAAAGGGGCAGGTGC  
 1819 TGGGCTCAGACCTGCCAAGAGCCATATCCGGGAGGACCTGCCCTGACCTAAGCCCCACCCAAAGGCCAACTCTCCAG  
 15 1899 TCCCTCAGCTCGGACACCTTCTCTCTCCAGATTCCAGTAACTCCCAATCTTCTCTGACAGGCCCAATCTTGTGAC  
 E P K S C D  
 1979 AAAACTCACACATGCCACCGTGCCAGGTAAGCCAGCCCAGGCTCGCCCTCCAGCTCAAGCGGGACAGGTGCCCTAG  
 K T H T C P P C P  
 2059 AGTAGCCTGCATCCAGGGACAGGCCCCAGCGGGTGTGACACGTCCACCTCCATCTCTTCTCAGCACCTGAACCTCTG  
 A P E L L  
 2139 GGGGACCGTCACTCTTCTTCCCCCAAAACCCAGGACACCCCTCATGATCTCCCGGACCCCTGAGGTACATGCGT  
 G G P S V F L P P K P K D T L M I S R T P E V T C V  
 2219 GGTGGTGACGTGAGCCACGAAGACCTGAGGTCAAGTCAACTGGTACGTGGACGGCGTGGAGTGCATAATGCCAAGA  
 V V D V S H E D P E V K F N W Y V D G V E V H N A K  
 25 2299 CAAAGCCCGGGAGGAGCAGTACAACAGCAGTACCGTGTGGTTCAGCGTCTCACCCTGTCACAGGACTGGCTGAAT  
 T K P R E E Q Y N S T Y R V V S V L T V L H Q D W L N  
 2379 GGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGGTGG  
 G K E Y K C K V S N K A L P A P I E K T I S K A K  
 2459 GACCCGTGGGGTGGAGGGCCACATGGACAGAGGCCGGCTCGGCCACCCCTCTGCCCTGAGAGTGACCGCTGTACCAACC  
 30 2539 TCTGTCCCTACAGGCGAGCCCCGAGAACCAGGTGTACACCTGCCCCCATCCCGGGATGAGCTGACCAAGAACAGGT  
 G Q P R E P Q V Y T L P P S R D E L T K N Q V  
 2619 CAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA  
 S L T C L V K G F Y P S D I A V E W E S N G Q P E N  
 2699 ACTACAAGACCACGCTCCCGTGTGGACTCCGACGGTCTCTTCTTCTCTACAGCAAGCTACCCGTGGACAGAGCAGG  
 35 N Y K T T P P V L D S D G S F L Y S K L T V D K S R  
 2779 TGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACAGCAGAGAGCCCTCTCCCT  
 W Q Q G N V F S C S V M H E A L H N H Y T Q K S L S L  
 2859 GTCTCCGGGTAAATGAGTGCGACGGCCGGCAAGCCCCCGCTCCCCGGGCTCTCGCGGTGCGACGAGGATGCTTGGCACGT  
 S P G K •  
 40 2939 ACCCCCTGTACATACTTCCCGGGCGCCAGCATGGAATAAA GCACCCAGCGCTGCCCTGGGCCCCCTGCGAGACTGTGAT  
 3019 GGTTCCTTCCACGGGTGAGGCGAGTCTGAGGCTGAGTGGCATGAGGGAGGAGAGCGGGTCCCACTGTCCCCCACTG  
 3099 GCCCAGGCTGTGACAGGTGTGCTGGGCGGCTAGGGTGGGGCTCAGCCAGGGGCTGCCCTCGGCAGGGTGGGGGATTTC  
 3179 CAGCGTGGCCCTCCCTCCAGCAGACCTGCCCCGGGCTGGGCCACGGGAAGCCCTAGGAGCCCTGGGGACAGACACA  
 3259 GCCCCGTGCTCTGTAGGAGACTGTCTGTTCTGTGAGCGCCCTGTCCTCCGACCTCCATGCCCACTCGGGGGATGCCCTA  
 45 3339 GTCCATGTGCGTAGGGACAGGCCCTCCCTACCCATCTACCCCCACGGCACTAACCCCTGGCTGCCCTGCCAGCCTCGC  
 3419 ACCCGCATGGGGACACAACCGACTCCGGGGACATGCACTCTCGGGCCCTGTGGAGGGAGTGGTGAGATGCCACACACA  
 3499 CACTCAGCCCAGACCGTTCAACAAACCCGCACTGAGGTTGGCCGGCCACAGGCCACACACACAGTCGACGCCT  
 3579 CACACACGGAGCCTCACCCGGGCGAACTGCACAGCACCAGACCAGAGCAAGGTCTCGCACAGTGAACACTCCTCGGA  
 3659 CACAGGCCCCACAGCCCCACGCGGCACCTCAAGGCCACAGGCTCTCGGCAGCTTCTCCACATGCTGACCTGCTCAG  
 50 3739 ACAAAACCCAGCCCTCCTCTCAAGGGTGCCCTGCAGCGCCACACACACAGGGGATCACACACACAGTCAAGTCC  
 3819 TGGCCCTGGCCCACTTCCAGTGCCGCCCTTCCCTGCAGGATCC (SEQ ID NO:16)

Human constant region DNA sequences can be isolated in accordance with well  
 known procedures from a variety of human cells, but preferably from immortalized B-  
 55 cells. Suitable source cells for the polynucleotide sequences and host cells for  
 immunoglobulin expression and secretion can be obtained from a number of sources well-  
 known in the art.

In addition to the humanized immunoglobulins specifically described herein, other  
 "substantially homologous" modified immunoglobulins can be readily designed and

manufactured utilizing various recombinant DNA techniques well known to those skilled in the art. For example, the framework regions can vary from the native sequences at the primary structure level by several amino acid substitutions, terminal and intermediate additions and deletions, and the like. Moreover, a variety of different human framework regions may be used singly or in combination as a basis for the humanized immunoglobulins of the present invention. In general, modifications of the genes may be readily accomplished by a variety of well-known techniques, such as site-directed mutagenesis.

Alternatively, polypeptide fragments comprising only a portion of the primary antibody structure may be produced, which fragments possess one or more immunoglobulin activities (e.g., complement fixation activity). These polypeptide fragments may be produced by proteolytic cleavage of intact antibodies by methods well known in the art, or by inserting stop codons at the desired locations in vectors using site-directed mutagenesis, such as after CH1 to produce Fab fragments or after the hinge region to produce F(ab')<sub>2</sub> fragments. Single chain antibodies may be produced by joining VL and VH with a DNA linker.

As stated previously, the polynucleotides will be expressed in hosts after the sequences have been operably linked to (i.e., positioned to ensure the functioning of) an expression control sequence. These expression vectors are typically replicable in the host organisms either as episomes or as an integral part of the host chromosomal DNA. Commonly, expression vectors will contain selection markers, e.g., tetracycline or neomycin, to permit detection of those cells transformed with the desired DNA sequences.

*E. coli* is a prokaryotic host useful particularly for cloning the polynucleotides of the present invention. Other microbial hosts suitable for use include bacilli, such as *Bacillus subtilis*, and other enterobacteriaceae, such as *Salmonella*, *Serratia*, and various *Pseudomonas* species. In these prokaryotic hosts, one can also make expression vectors, which will typically contain expression control sequences compatible with the host cell (e.g., an origin of replication). In addition, any of a number of well-known promoters may be present, such as the lactose promoter system, a tryptophan (trp) promoter system, a beta-lactamase promoter system, or a promoter system from phage lambda. The promoters will typically control expression, optionally with an operator sequence, and

have ribosome binding site sequences and the like, for initiating and completing transcription and translation.

Other microbes, such as yeast, may also be used for expression. *Saccharomyces* is a preferred host, with suitable vectors having expression control sequences, such as  
5 promoters, including 3-phosphoglycerate kinase or other glycolytic enzymes, and an origin of replication, termination sequences and the like as desired.

In addition to microorganisms, mammalian tissue cell culture may also be used to express and produce the polypeptides of the present invention. Eukaryotic cells are actually preferred, because a number of suitable host cell lines capable of secreting intact  
10 immunoglobulins have been developed in the art, and include the CHO cell lines, various COS cell lines, Syrian Hamster Ovary cell lines, HeLa cells, preferably myeloma cell lines, transformed B-cells, human embryonic kidney cell lines, or hybridomas. Expression vectors for these cells can include expression control sequences, such as an origin of replication, a promoter, an enhancer, and necessary processing information sites,  
15 such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. Preferred expression control sequences are promoters derived from immunoglobulin genes, SV40, Adenovirus, Bovine Papilloma Virus, cytomegalovirus and the like.

The vectors containing the polynucleotide sequences of interest (e.g., the heavy  
20 and light chain encoding sequences and expression control sequences) can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment or electroporation may be used for other cellular hosts.

25 Once expressed, the antibodies can be purified according to standard procedures, including ammonium sulfate precipitation, ion exchange, *affinity*, *reverse phase*, hydrophobic interaction column chromatography, gel electrophoresis, and the like. Substantially pure immunoglobulins of at least about 90 to 95% homogeneity are preferred, and 98 to 99% or more homogeneity most preferred, for pharmaceutical uses.  
30 Once purified, partially or to homogeneity as desired, the polypeptides may then be used therapeutically or prophylactically, as directed herein.

The antibodies (including immunologically reactive fragments) are administered to a subject at risk for or exhibiting A $\beta$ -related symptoms or pathology such as clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical amyloid angiopathy, using standard administration techniques, preferably peripherally (*i.e.* not by administration into the central nervous system) by intravenous, intraperitoneal, subcutaneous, pulmonary, transdermal, intramuscular, intranasal, buccal, sublingual, or suppository administration. Although the antibodies may be administered directly into the ventricular system, spinal fluid, or brain parenchyma, and techniques for addressing these locations are well known in the art, it is not necessary to utilize these more difficult procedures. The antibodies of the invention are effective when administered by the more simple techniques that rely on the peripheral circulation system. The advantages of the present invention include the ability of the antibody to exert its beneficial effects even though not provided directly to the central nervous system itself. Indeed, it has been demonstrated that the amount of antibody that crosses the blood-brain barrier is  $\leq 0.1\%$  of plasma levels.

The pharmaceutical compositions for administration are designed to be appropriate for the selected mode of administration, and pharmaceutically acceptable excipients such as, buffers, surfactants, preservatives, solubilizing agents, isotonicity agents, stabilizing agents and the like are used as appropriate. Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton PA, latest edition, incorporated herein by reference, provides a compendium of formulation techniques as are generally known to practitioners.

The concentration of the humanized antibody in formulations may range from as low as about 0.1% to as much as 15 or 20% by weight and will be selected primarily based on fluid volumes, viscosities, and so forth, in accordance with the particular mode of administration selected. Thus, a pharmaceutical composition for injection could be made up to contain in 1 mL of phosphate buffered saline from 1 to 100 mg of the humanized antibody of the present invention. The formulation could be sterile filtered after making the formulation, or otherwise made microbiologically acceptable. A typical composition for intravenous infusion could have a volume as much as 250 mL of fluid, such as sterile Ringer's solution, and 1-100 mg per mL, or more in antibody concentration. Therapeutic agents of the invention can be frozen or lyophilized for storage and

reconstituted in a suitable sterile carrier prior to use. Lyophilization and reconstitution can lead to varying degrees of antibody activity loss (e.g. with conventional immune globulins, IgM antibodies tend to have greater activity loss than IgG antibodies). Dosages may have to be adjusted to compensate. The pH of the formulation will be selected to  
5 balance antibody stability (chemical and physical) and comfort to the patient when administered. Generally, pH between 4 and 8 is tolerated.

Although the foregoing methods appear the most convenient and most appropriate for administration of proteins such as humanized antibodies, by suitable adaptation, other techniques for administration, such as transdermal administration and oral administration  
10 may be employed provided proper formulation is designed. In addition, it may be desirable to employ controlled release formulations using biodegradable films and matrices, or osmotic mini-pumps, or delivery systems based on dextran beads, alginate, or collagen. In summary, formulations are available for administering the antibodies of the invention and are well-known in the art and may be chosen from a variety of options.  
15 Typical dosage levels can be optimized using standard clinical techniques and will be dependent on the mode of administration and the condition of the patient.

The following examples are intended to illustrate but not to limit the invention. Because the examples here describe experiments conducted in murine systems, the use of murine monoclonal antibodies is satisfactory. However, in the treatment methods of the  
20 invention intended for human use, humanized forms of the antibodies with the immunospecificity corresponding to that of antibody 3D6 are preferred.

### Example 1

#### Synthesis of Humanized Antibody 3D6

25 Cells and antibodies. Mouse myeloma cell line Sp2/0 was obtained from ATCC (Manassas, VA) and maintained in DME medium containing 10% FBS (Cat # SH30071.03, HyClone, Logan, UT) in a 37°C CO<sub>2</sub> incubator. Mouse 3D6 hybridoma cells were first grown in RPMI-1640 medium containing 10% FBS (HyClone), 10 mM HEPES, 2 mM glutamine, 0.1 mM non-essential amino acids, 1 mM sodium pyruvate, 25  
30 µg/ml gentamicin, and then expanded in serum-free media (Hybridoma SFM, Cat # 12045-076, Life Technologies, Rockville, MD) containing 2% low Ig FBS (Cat # 30151.03, HyClone) to a 1.5 liter volume in roller bottles. Mouse monoclonal antibody

3D6 (Mu3D6) was purified from the culture supernatant by affinity chromatography using a protein-G Sepharose column. Biotinylated Mu3D6 was prepared using EZ-Link Sulfo-NHS-LC-LC-Biotin (Cat # 21338ZZ, Pierce, Rockford, IL).

Cloning of variable region cDNAs. Total RNA was extracted from approximately 5 10<sup>7</sup> hybridoma cells using TRIzol reagent (Cat. # 15596-026 Life Technologies) and poly(A)<sup>+</sup> RNA was isolated with the PolyAtract mRNA Isolation System (Cat. # Z5310, Promega, Madison, WI) according to the suppliers' protocols. Double-stranded cDNA was synthesized using the SMART<sup>TM</sup>RACE cDNA Amplification Kit (Cat. # K1811-1, Clontech, Palo Alto, CA) following the supplier's protocol. The variable region cDNAs 10 for the light and heavy chains were amplified by polymerase chain reaction (PCR) using 3' primers that anneal respectively to the mouse kappa and gamma chain constant regions, and a 5' universal primer provided in the SMART<sup>TM</sup>RACE cDNA Amplification Kit. For VL PCR, the 3' primer has the sequence:

15 5' -TATAGAGCTCAAGCTTGGATGGTGGGAAGATGGATACAGTTGGTGC-3'  
[SEQ ID NO:13]

with residues 17- 46 hybridizing to the mouse Ck region. For VH PCR, the 3' primers have the degenerate sequences:

20

5' -TATAGAGCTCAAGCTTCCAGTGGATAGACCGATGGGGCTGTCGTTTTGGC-3'

T

[SEQ ID NO:14]

25 with residues 17 - 50 hybridizing to mouse gamma chain CH1. The VL and VH cDNAs were subcloned into pCR4Blunt-TOPO vector (Cat. # 45-0031, Invitrogen, Carlsbad, CA) for sequence determination. DNA sequencing was carried out by PCR cycle sequencing reactions with fluorescent dideoxy chain terminators (Applied Biosystems, Foster City, CA) according to the manufacturer's instructions. The sequencing reactions were  
30 analyzed on a Model 377 DNA Sequencer (Applied Biosystems).

Construction of humanized 3D6 (Hu3D6) variable regions. Humanization of the mouse antibody V regions was carried out as outlined by Queen et al., (1989), *op. cit.* The human V region framework used as acceptor for Mu3D6 CDRs was chosen based on

sequence homology. The computer programs ABMOD and ENCAD [Levitt, M., J. Mol. Biol. 168:595-620 (1983)] were used to construct a molecular model of the variable regions. Amino acids in the humanized V regions that were predicted to have contact with CDRs were substituted with the corresponding residues of Mu3D6. This was done  
5 at residues 49, 73, and 98 in the heavy chain and at residue 41 in the light chain. The amino acids in the humanized V region that were found to be rare in the same V-region subgroup were changed to the consensus amino acids to eliminate potential immunogenicity. This was done at residues 6 and 91 in the heavy chain.

The light and heavy chain variable region genes were constructed and amplified  
10 using eight overlapping synthetic oligonucleotides ranging in length from approximately 65 to 80 bases [He, X. Y., et al., J. Immunol. 160: 029-1035 (1998)]. The oligonucleotides were annealed pairwise and extended with the Klenow fragment of DNA polymerase I, yielding four double-stranded fragments. The resulting fragments were denatured, annealed pairwise, and extended with Klenow, yielding two fragments. These  
15 fragments were denatured, annealed pairwise, and extended once again, yielding a full-length gene. The resulting product was amplified by PCR using the Expand High Fidelity PCR System (Cat. # 1 732 650, Roche Molecular Biochemicals, Indianapolis, IN). The PCR-amplified fragments were gel-purified and cloned into pCR4Blunt-TOPO vector. After sequence confirmation, the VL and VH genes were digested with MluI and XbaI,  
20 gel-purified, and subcloned respectively into vectors for expression of light and heavy chains to make pVk-Hu3D6 and pVg1-Hu3D6 [Co, M. S., et al., J. Immunol. 148:1149-1154 (1992)]. The mature humanized 3D6 antibody expressed from these plasmids has the light chain of SEQ ID NO:11 and the heavy chain of SEQ ID NO:12.

Stable transfection. Stable transfection into mouse myeloma cell line Sp2/0 was  
25 accomplished by electroporation using a Gene Pulser apparatus (BioRad, Hercules, CA) at 360 V and 25  $\mu$ F as described (Co, et al., 1992, *op. cit.*). Before transfection, pVk-Hu3D6 and pVg1-Hu3D6 plasmid DNAs were linearized using FspI and BstZ171, respectively. Approximately  $10^7$  Sp2/0 cells were transfected with 20  $\mu$ g of pVk-Hu3D6 and 40  $\mu$ g of pVg1-Hu3D6. The transfected cells were suspended in DME medium  
30 containing 10% FBS and plated into several 96-well plates. After 48 hr, selection media (DME medium containing 10% FBS, HT media supplement, 0.3 mg/ml xanthine and 1  $\mu$ g/ml mycophenolic acid) was applied. Approximately 10 days after the initiation of the

selection, culture supernatants were assayed for antibody production by ELISA as shown below. High yielding clones were expanded in DME medium containing 10% FBS and further analyzed for antibody expression. Selected clones were then adapted to growth in Hybridoma SFM.

5        Measurement of antibody expression by ELISA. Wells of a 96-well ELISA plate (Nunc-Immuno plate, Cat # 439454, NalgeNunc, Naperville, IL) were coated with 100  $\mu$ l of 1  $\mu$ g/ml goat anti-human IgG, Fc  $\gamma$  fragment specific, polyclonal antibodies (Cat # 109-005-098, Jackson ImmunoResearch, West Grove, PA) in 0.2 M sodium carbonate-bicarbonate buffer (pH 9.4) overnight at 4°C. After washing with Washing Buffer (PBS  
10        containing 0.1% Tween 20), wells were blocked with 400  $\mu$ l of Superblock Blocking Buffer (Cat # 37535, Pierce) for 30 min and then washed with Washing Buffer. Samples containing Hu3D6 were appropriately diluted in ELISA Buffer (PBS containing 1% BSA and 0.1% Tween 20) and applied to ELISA plates (100  $\mu$ l per well). As a standard, humanized anti-CD33 IgG1 monoclonal antibody HuM195 (Co, *et al.*, 1992, *op. cit.*) was  
15        used. The ELISA plate was incubated for 2 hr at room temperature and the wells were washed with Washing Buffer. Then, 100  $\mu$ l of 1/1,000-diluted HRP-conjugated goat anti-human kappa polyclonal antibodies (Cat # 1050-05, Southern Biotechnology, Birmingham, AL) in ELISA Buffer was applied to each well. After incubating for 1 hr at room temperature and washing with Washing Buffer, 100  $\mu$ l of ABTS substrate (Cat #s  
20        507602 and 506502, Kirkegaard and Perry Laboratories, Gaithersburg, MD) was added to each well. Color development was stopped by adding 100  $\mu$ l of 2% oxalic acid per well. Absorbance was read at 415 nm using an OPTImax microplate reader (Molecular Devices, Menlo Park, CA).

Purification of Hu3D6. One of the high Hu3D6-expressing Sp2/0 stable  
25        transfectants (clone #40) was adapted to growth in Hybridoma SFM and expanded to 2 liters in roller bottles. Spent culture supernatant was harvested when cell viability reached 10% or below and loaded onto a protein-A Sepharose column. The column was washed with PBS before the antibody was eluted with 0.1 M glycine-HCl (pH 2.5), 0.1 M NaCl. The eluted protein was dialyzed against 3 changes of 2 liters of PBS and filtered  
30        through a 0.2  $\mu$ m filter prior to storage at 4°C. Antibody concentration was determined by measuring absorbance at 280 nm (1 mg/ml = 1.4 A<sub>280</sub>). SDS-PAGE in Tris-glycine buffer was performed according to standard procedures on a 4-20% gradient gel (Cat #



EC6025, Novex, San Diego, CA). Purified humanized 3D6 antibody is reduced and run on an SDS- PAGE gel. The whole antibody shows two bands of approximate molecular weights 25 kDa and 50 kDa. These results are consistent with the molecular weights of the light chain and heavy chain, or with the molecular weight of the chain(s) comprising a  
5 fragment, calculated from their amino acid compositions.

## Example 2

### *In vitro* binding properties of humanized 3D6 antibody

The binding efficacy of humanized 3D6 antibody, synthesized and purified as  
10 described above, was compared with the mouse 3D6 antibody using biotinylated mouse 3D6 antibody in a comparative ELISA. Wells of a 96-well ELISA plate (Nunc-Immuno plate, Cat # 439454, NalgeNunc) were coated with 100  $\mu$ l of  $\beta$ -amyloid peptide (1-42) in 0.2 M sodium carbonate/bicarbonate buffer (pH 9.4) (0.3  $\mu$ g/mL) overnight at 4°C.

After washing the wells with phosphate buffered saline (PBS) containing 0.1%  
15 Tween 20 (Washing Buffer) using an ELISA plate washer, the wells were blocked by adding 300  $\mu$ L of SuperBlock reagent (Pierce) per well. After 30 minutes of blocking, the wells were washed with Washing Buffer and excess liquid was removed.

A mixture of biotinylated Mu3D6 (0.2  $\mu$ g/ml final concentration) and competitor antibody (Mu3D6 or Hu3D6; starting at 300  $\mu$ g/ml final concentration and serial 3-fold  
20 dilutions) in ELISA Buffer were added in triplicate in a final volume of 100  $\mu$ l per well. As a no-competitor control, 100  $\mu$ l of 0.2  $\mu$ g/ml biotinylated Mu3D6 was added. As a background control, 100  $\mu$ l of ELISA Buffer was added. The ELISA plate was incubated at room temperature for 90 min. After washing the wells with Washing Buffer, 100  $\mu$ l of 1  $\mu$ g/ml HRP-conjugated streptavidin (Cat # 21124, Pierce) was added to each well. The  
25 plate was incubated at room temperature for 30 min and washed with Washing Buffer. For color development, 100  $\mu$ l/well of ABTS Peroxidase Substrate (Kirkegaard & Perry Laboratories) was added. Color development was stopped by adding 100  $\mu$ l/well of 2% oxalic acid. Absorbance was read at 415 nm. The absorbances were plotted against the log of the competitor concentration, curves were fit to the data points (using Prism) and  
30 the IC50 was determined for each antibody using methods well-known in the art.

The mean IC50 for mouse 3D6 was 2.7  $\mu$ g/mL (three separate experiments, standard deviation = 0.6  $\mu$ g/mL) and for humanized 3D6 was 3.3  $\mu$ g/mL (three separate

experiments, standard deviation = 0.8  $\mu\text{g/mL}$ ). A second set of three experiments was carried out, essentially as described above, and the mean  $\text{IC}_{50}$  for mouse 3D6 was determined to be 3.97  $\mu\text{g/mL}$  (SD = 0.15  $\mu\text{g/mL}$ ) and for humanized 3D6, the  $\text{IC}_{50}$  was determined to be 3.97  $\mu\text{g/mL}$  (SD = 0.20  $\mu\text{g/mL}$ ). On the basis of these results, we  
5 conclude that humanized 3D6 has binding properties that are very similar to those of the mouse antibody 3D6. Therefore, we expect that humanized 3D6 has very similar *in vitro* and *in vivo* activities compared with mouse 3D6 and will exhibit in humans the same effects demonstrated with mouse 3D6 in mice.

### Example 3

#### *In vitro* binding properties of mouse and humanized antibodies 3D6

Antibody affinity ( $\text{KD} = \text{Kd} / \text{Ka}$ ) was determined using a BIAcore biosensor 2000 and data analyzed with BIAevaluation (v. 3.1) software. A capture antibody (rabbit anti-mouse or anti-human IgG) was coupled via free amine groups to carboxyl groups on flow  
15 cell 2 of a biosensor chip (CM5) using N-ethyl-N-dimethylaminopropyl carbodiimide and N-hydroxysuccinimide (EDC/NHS). A non-specific rabbit IgG was coupled to flow cell 1 as a background control. Monoclonal antibodies were captured to yield 300 resonance units (RU). Amyloid-beta 1-40 or 1-42 (Biosource International, Inc.) was then flowed over the chip at decreasing concentrations (1000 to 0.1 times KD). To regenerate the  
20 chip, bound anti-A $\beta$  antibody was eluted from the chip using a wash with glycine-HCl (pH 2). A control injection containing no amyloid-beta served as a control for baseline subtraction. Sensorgrams demonstrating association and dissociation phases were analyzed to determine Kd and Ka. The affinity (KD) of mouse antibody 3D6 for A $\beta$  1-42 was determined to be 2.4 nM, and the affinity of humanized 3D6, prepared essentially as  
25 described in Example 1, was determined to be 2.3 nM.

### Example 4

#### Epitope mapping of mouse and humanized 3D6

The BIAcore is an automated biosensor system for measuring molecular  
30 interactions [Karlsson R., *et al. J. Immunol. Methods* 145:229-240 (1991)]. The advantage of the BIAcore over other binding assays is that binding of the antigen can be measured without having to label or immobilize the antigen (i.e. the antigen maintains a

more native conformation). The BIAcore methodology was used to assess the binding of various amyloid-beta peptide fragments to either mouse 3D6 or humanized 3D6 (prepared substantially as described in Example 1). All dilutions were made with HEPES buffered saline containing Tween 20. A single concentration of a variety of fragments of human A $\beta$  or mouse A $\beta$  1-40 (BioSource International) was used. Human amyloid beta fragments 1-10 and 1-20 bound to mouse 3D6 and to humanized 3D6, while human A $\beta$  fragments 10-20 and 16-25 did not bind to either antibody. Neither mouse 3D6 nor humanized 3D6 bound mouse A $\beta$  1-40. Using this methodology, the binding epitope for both mouse and humanized 3D6 appears to be between amino acids 1 and 10 of human A $\beta$ .

#### Example 5

##### Effects of administration of 3D6

Unless otherwise stated, all studies used APP<sup>V717F</sup> (PDAPP) transgenic mice, and all injections were i.p. In general, a control group of mice received injections of saline.

Six weeks of weekly injection of 360  $\mu$ g of 3D6 in old, hemizygous mice (24 month) lowered hippocampal insoluble A $\beta_{\text{total}}$  by 10% and A $\beta$  1-42 by 1% (N.S., not statistically significant) in 9 animals per control group and 10 animals per antibody group. In the cortex, mean insoluble A $\beta_{\text{total}}$  was lower by 33% and A $\beta$  1-42 by 47% ( $p < 0.05$ ), while insoluble A $\beta$  1-40 increased by 100%.

In hemizygous, 4 month old mice, administration of 360  $\mu$ g of 3D6 per animal: 1) raised average plasma A $\beta$  1-40 and A $\beta$  1-42 levels approximately 6-fold and 9-fold, respectively, by 24 hours after administration; and 2) had no significant effect on soluble A $\beta$  1-40 in the cortex after 24 hours compared with saline control (5 animals per group). In another study with hemizygous, 3 month old mice, administration of 360  $\mu$ g of 3D6 per animal raised average plasma A $\beta$  1-42 levels approximately 8-fold by 24 hours after administration.

Administration of 360  $\mu$ g of 3D6 per animal (5 animals per group, saline control): raised average plasma A $\beta$  1-40 and A $\beta$  1-42 levels approximately 92-fold and 32-fold, respectively, by 24 hours after administration ( $p < 0.05$ ); lowered cortical insoluble A $\beta$  1-40 by 42% ( $p < 0.05$ ) and A $\beta$  1-42 by 27% (N.S.), but increased A $\beta_{\text{total}}$  by 35% (N.S.); had no consistent or significant effect on soluble or insoluble A $\beta$  1-40, A $\beta$  1-42, or A $\beta_{\text{total}}$  in

the hippocampus after 24 hours; in the cerebellum, increased soluble A $\beta$  1-42 by 80% ( $p < 0.001$ ) and A $\beta_{\text{total}}$  by 68% (N.S.), but lowered soluble A $\beta$  1-40 by 6% (N.S.); and in the cerebellum, lowered insoluble A $\beta$  1-40, A $\beta$  1-42, and A $\beta_{\text{total}}$  by 35% ( $p < 0.01$ ), 21% (N.S.), and 12% (N.S.), respectively.

- 5 In young mice, administration of 360  $\mu\text{g}$  of 3D6 per animal (5 per group): 1) raised average plasma A $\beta$  1-42 levels approximately 3-fold by 24 hours after administration; and 2) in the cortex, lowered insoluble A $\beta$  1-40 about 10% and increased insoluble A $\beta$  1-42 about 12%.

- Studies were conducted to assess the effects of 3D6 on formation of stable  
10 A $\beta$ :antibody complexes in biological fluids, plasma A $\beta$  concentrations acutely after administration, cognitive performance after acute or chronic administration, and guanidine-extracted and immunohistochemically-detected A $\beta$  deposition (in brain) after chronic administration.

- Mice (3 months of age) were injected with 360  $\mu\text{g}$  of 3D6. Twenty-four hours  
15 following antibody administration plasma was collected and proteins were resolved by gel electrophoresis under native (non-denaturing conditions) on a polyacrylamide gel. Following transfer of size fractionated proteins to a solid matrix, complexes were immunodetected with biotinylated antibody and visualized with enhanced chemiluminescence. Unlike certain other anti-A $\beta$  antibodies, no complex was detected  
20 with 3D6.

- Young (2-3 months of age) mice were injected with 3D6. At various times following antibody administration, plasma was collected and various A $\beta$  species were determined by a sandwich ELISA. Administration of 3D6 resulted in a dose- and time-dependent increase in plasma A $\beta$  levels. A $\beta_{1-40}$  levels increased to a greater degree than  
25 A $\beta_{1-42}$  levels following 3D6 administration. In an additional study, young APP<sup>V717F</sup> tg mice were treated with 360  $\mu\text{g}$  3D6 and plasma A $\beta$  levels were measured at 0.5, 3, 6, and 24 h following injection. 3D6 increased plasma A $\beta$  levels in a time-dependent manner.

- Extensive behavioral characterization of APP<sup>V717F</sup> tg mice has been performed using several memory paradigms (bar-press, 8 arm-radial maze, object recognition).  
30 These mice are impaired in several learning and memory tasks, and deficits in the object recognition (OR) task worsen with age. Therefore, the OR task has been used to assess

learning and memory in APP<sup>V717F</sup> tg mice. Performance in the OR task is preferentially dependent on the integrity of the medial temporal lobe (perirhinal and entorhinal cortices). The OR test relies on the spontaneous tendency of rodents to preferentially explore a novel versus familiar object.

5        On the first day of testing, mice were allowed to habituate to an open field chamber for 50 minutes. The following day, mice were placed back into the open field for two 10-min trials. During trial one, mice were allowed to explore the open field in the presence of an object (e.g., marble or die). Following a 3-hr inter-trial delay, mice were placed back into the open field with the familiar object (the same object explored  
10       previously during trial 1) as well as a novel object. The time spent exploring the novel object as well as the familiar object was recorded and a recognition index (the ratio of time spent exploring the novel object x 100/ total time spent exploring both objects) was calculated for each mouse. Administration of 360 µg of 3D6 per animal 24 hours prior to the habituation session in 11-12 month old APP<sup>V717F</sup> tg mice improved OR performance  
15       in 2 of 8 mice tested (p<0.05).

         Homozygous tg mice (5-6 months old) were administered weekly injections of PBS and 72, 217, and 360 µg of a non-specific IgG or 3D6 (n = 19-30) for 5 months. At necropsy, the brains were removed and processed for Aβ ELISA assays and immunohistochemical analysis of parenchymal Aβ burden. Cortical and hippocampal  
20       tissues were homogenized in PBS. PBS-insoluble Aβ was subsequently extracted from the pellets by homogenization in 5.5 M guanidine-HCl. Following homogenization, the samples were nutated for at least 24 h prior to centrifugation and collection of the guanidine extract. PBS-soluble and guanidine-extracted tissue preparations were stored at -80°C for subsequent Aβ ELISA determinations. Immunohistochemical (IHC) analysis  
25       of parenchymal Aβ burden was carried out as follows. Eight (8) µm paraffin embedded paraformaldehyde fixed tissues were labeled with rabbit polyclonal anti-Aβ antibody (against Aβ 15-30) and followed by anti-rabbit IgG fluorescent detection. Eight (8) sections of brain (7 IHC, 1 control) were examined from each animal. Treatment with 3D6 (360 µg) markedly and significantly reduced cortical guanidine-extracted Aβ1-42 (by  
30       ELISA) and cortical and hippocampal Aβ plaque burden (by IHC), but no effect was observed at lower 3D6 doses. Although no effect on guanidine extracted Aβ1-42 was

observed at lower 3D6 doses, these doses significantly reduced cortical and hippocampal A $\beta$  plaque burden (by IHC).

Radiolabeled (15  $\mu$ Ci/mouse, 0.5 mg/mouse) 3D6 was administered to ICR (non-transgenic) mice in order to evaluate kinetics and brain distribution of the antibody after administration by the intravenous route. Plasma kinetics for 3D6 immunoreactivity demonstrated a half-life of elimination of approximately 5 days. TCA-precipitable radioactivity was greater than 95% of the total plasma counts throughout the study, and declined in the plasma compartment with a terminal half-life of 3-4 days. The observation that plasma radioactivity remained predominantly TCA-precipitable throughout the study suggests that the radiolabeled antibody was not significantly proteolytically degraded, nor was the 125-I label cleaved from the antibody over the time course studied. The shapes of the concentration versus time profiles as measured by ELISA and radioactivity were generally similar, with some differences in the terminal phases. There was no apparent accumulation of radiolabel in any tissue, including brain. Distribution of radioactivity to the brain was minimal. The amount of radioactivity associated with the brain samples in this experiment cannot be clearly distinguished from contamination by the blood compartment during tissue processing or from antibody associated with endothelial cells in the brain vasculature.

Nine month old, hemizygous mice received PBS, a non-specific IgG, or 3D6 (500  $\mu$ g/week) by weekly injection for six months (PBS, 11 animals; IgG, 13 animals; and 3D6, 14 animals). Weak, but statistically significant, A $\beta$  lowering in the cortex (compared to IgG) and hippocampus (compared to IgG or combined PBS/IgG controls) was seen. Immunohistochemical (IHC) analysis showed strong reductions in A $\beta$  plaque burden in the cortex and hippocampus of 3D6-treated mice (94% and 85% reductions, respectively, versus PBS control;  $p < 0.05$ , and  $p < 0.01$ , respectively).

### Example 6

#### Administration of humanized 3D6

A preparation of an anti-A $\beta$  antibody comprising a light chain having the amino acid sequence of SEQ ID NO:11 and a heavy chain having the amino acid sequence of SEQ ID NO:12 (a humanized 3D6) was administered as a single intravenous bolus injection to two groups of 12 male marmosets at doses of 1 and 10 mg/kg.

Concentrations of immunoreactive anti-A $\beta$  antibody declined with a half-life or elimination of approximately 4 days.  $C_{max}$  and AUC parameters increased proportionally between the 1 and 10 mg/kg dose levels. The administration of humanized 3D6 to marmosets resulted in 18 or 29-fold increase in plasma A $\beta_{1-40}$  immunoreactivity after 8 hours, compared with predose concentrations in the 1 and 10 mg/kg dose groups, respectively. Animals at both dose levels had concentrations of A $\beta_{1-40}$  immunoreactivity above baseline levels up to 2 weeks after antibody administration. Kinetic analysis of concentrations of A $\beta_{1-40}$  immunoreactivity showed that the half-life of elimination of A $\beta_{1-40}$  immunoreactivity was comparable to that of the antibody (~4 days). The pharmacokinetics of humanized 3D6 were also evaluated in male *cynomolgus* monkeys after a single intravenous administration of 1 mg/kg. Analysis of immunoreactivity showed that humanized 3D6 was eliminated from the plasma with a half-life of approximately 11-12 days.

We claim:

1. Humanized 3D6 antibody.

2. A humanized antibody, or fragment thereof, comprising a humanized light chain comprising three light chain complementarity determining regions (CDRs) from the mouse monoclonal antibody 3D6 and a light chain variable region framework sequence  
5 from a human immunoglobulin light chain; and a humanized heavy chain comprising three heavy chain CDRs from the mouse monoclonal antibody 3D6 and a heavy chain variable region framework sequence from a human immunoglobulin heavy chain; wherein the light chain CDRs have the following amino acid sequences:

10 light chain CDR1:

1 5 10 15  
Lys Ser Ser Gln Ser Leu Leu Asp Ser Asp Gly Lys Thr Tyr Leu Asn  
(SEQ ID NO:1)

15 light chain CDR2:

1 5  
Leu Val Ser Lys Leu Asp Ser (SEQ ID NO:2)

light chain CDR3:

20 1  
Trp Gln Gly Thr His Phe Pro Arg Thr (SEQ ID NO:3)

and the heavy chain CDRs have the following amino acid sequences:

heavy chain CDR1:

25 1 5  
Asn Tyr Gly Met Ser (SEQ ID NO:4)

heavy chain CDR2:

30 1 5 10 15  
Ser Ile Arg Ser Gly Gly Gly Arg Thr Tyr Tyr Ser Asp Asn Val Lys Gly  
(SEQ ID NO:5)

and, heavy chain CDR3:

35 1 5 10  
Tyr Asp His Tyr Ser Gly Ser Ser Asp Tyr (SEQ ID NO:6).

3. A humanized antibody or fragment thereof comprising a humanized light chain variable region having the sequence of SEQ ID NO:7 and a humanized heavy variable region having the sequence of SEQ ID NO:8.



4. The humanized antibody or fragment thereof of claim 3 having a light chain variable region of the sequence given by SEQ ID NO:9 and a heavy chain variable region given by SEQ ID NO:10.
5. The humanized antibody or fragment thereof of claim 3 having a light chain of the sequence given by SEQ ID NO:11 and a heavy chain of the sequence given by SEQ ID NO:12.
6. An antibody fragment obtainable by enzymatic cleavage of the humanized antibody of any one of claims 1 - 5.
7. An Fab or F(ab')<sub>2</sub> fragment of any one of the humanized antibodies of claims 1 - 5.
8. The F(ab')<sub>2</sub> fragment of claim 7.
9. The Fab fragment of claim 7.
10. The humanized antibody or fragment of any one of claims 1 - 9, which is a single chain antibody.
11. The humanized antibody or fragment of any one of claims 1 - 10 that is an IgG<sub>1</sub> immunoglobulin isotype.
12. The humanized antibody or fragment of any one of claims 1 - 11, wherein the antibody or fragment thereof is produced in a host cell selected from the group consisting of a myeloma cell, a chinese hamster ovary cell, a syrian hamster ovary cell, and a human embryonic kidney cell.
13. A polynucleotide compound, comprising a sequence coding for the light chain or the heavy chain of the humanized antibody of any one of claims 1 - 12, or a fragment thereof.

14. A polynucleotide sequence, which when expressed in a suitable host cell, yields an antibody of any one of claims 1 – 12.

15. The polynucleotide of claim 13 or 14 selected from the group consisting of SEQ ID NO: 15, SEQ ID NO: 17, and a polynucleotide comprising a sequence that codes  
5 for the light chain variable region given by SEQ ID NO:7, SEQ ID NO:9, or SEQ ID NO: 11.

16. The polynucleotide of claim 13 or 14 selected from the group consisting of SEQ ID NO:16, SEQ ID NO:18, and a polynucleotide comprising a sequence that codes for the heavy chain variable region given by SEQ ID NO:8, SEQ ID NO:10, or SEQ ID  
10 NO:12.

17. An expression vector for expressing the antibody of any one of claims 1 - 12 comprising the polynucleotide sequence of any one of claims 13 - 16.

18. A cell transfected with the expression vector of claim 17.

19. A cell transfected with two expression vectors of claim 17, wherein a first  
15 vector comprises the polynucleotide sequence coding for the light chain and a second vector comprises the sequence coding for the heavy chain.

20. A cell that is capable of expressing the humanized antibody or fragment of any one of claims 1 – 12.

21. The cell of any one of claims 18 – 20, wherein the cell is selected from the  
20 group consisting of a myeloma cell, a chinese hamster ovary cell, a syrian hamster ovary cell, and a human embryonic kidney cell.

22. A pharmaceutical composition comprising the humanized antibody or fragment of any one of claims 1 – 12, and a pharmaceutically acceptable excipient.

23. A method of treating Down's syndrome, clinical or pre-clinical Alzheimer's disease, or clinical or pre-clinical cerebral amyloid angiopathy in a human subject, comprising administering to the human subject an effective amount of a humanized antibody or fragment of any one of claims 1 – 12.

5           24. A method to inhibit the formation of A $\beta$  plaque in the brain of a human subject, comprising administering to the human subject an effective amount of the humanized antibody or fragment of any one of claims 1 – 12.

25. A method to reduce A $\beta$  plaque in the brain of a human subject, comprising administering to the human subject an effective amount of a humanized antibody or  
10 fragment of any one of claims 1 – 12.

26. The method of either of claims 24 – 25, wherein the subject is diagnosed with clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical cerebral amyloid angiopathy.

27. The method of any one of claims 24 – 25, wherein the subject is not  
15 diagnosed with clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical cerebral amyloid angiopathy.

28. Use of the humanized antibody or a fragment thereof according to any one of Claims 1 – 12 for the manufacture of a medicament, including prolonged expression of recombinant sequences of the antibody or antibody fragment in human tissues, for treating  
20 clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical cerebral amyloid angiopathy.

29. Use of the humanized antibody or fragment of any one of claims 1 – 12 for the manufacture of a medicament for treating Alzheimer's disease.

## SEQUENCE LISTING

<110> Eli Lilly and Company

<120> Humanized Antibodies

<130> X14958

<150> 60/287539

<151> 2001-04-30

<160> 20

<170> PatentIn version 3.1

<210> 1

<211> 16

<212> PRT

<213> Mus sp.

<400> 1

Lys Ser Ser Gln Ser Leu Leu Asp Ser Asp Gly Lys Thr Tyr Leu Asn  
1 5 10 15

<210> 2

<211> 7

<212> PRT

<213> mus sp.

<400> 2

Leu Val Ser Lys Leu Asp Ser  
1 5

<210> 3

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; mus sp.

&lt;400&gt; 3

Trp Gln Gly Thr His Phe Pro Arg Thr  
 1 5

&lt;210&gt; 4

&lt;211&gt; 5

&lt;212&gt; PRT

&lt;213&gt; mus sp.

&lt;400&gt; 4

Asn Tyr Gly Met Ser  
 1 5

&lt;210&gt; 5

&lt;211&gt; 17

&lt;212&gt; PRT

&lt;213&gt; mus sp.

&lt;400&gt; 5

Ser Ile Arg Ser Gly Gly Gly Arg Thr Tyr Tyr Ser Asp Asn Val Lys  
 1 5 10 15

Gly

&lt;210&gt; 6

&lt;211&gt; 10

&lt;212&gt; PRT

&lt;213&gt; mus sp.

&lt;400&gt; 6

Tyr Asp His Tyr Ser Gly Ser Ser Asp Tyr  
 1 5 10

<210> 7  
<211> 113  
<212> PRT  
<213> Artificial sequence

<220>  
<223> humanized antibody  
<220>  
<221> MISC\_FEATURE  
<222> (1)..(1)  
<223> Xaa=Asp or Tyr

<220>  
<221> MISC\_FEATURE  
<222> (7)..(7)  
<223> Xaa=Ser or Thr

<220>  
<221> MISC\_FEATURE  
<222> (10)..(10)  
<223> Xaa=Ser or Thr

<220>  
<221> MISC\_FEATURE  
<222> (15)..(15)  
<223> Xaa=Leu, Ile, or Val

<220>  
<221> MISC\_FEATURE  
<222> (50)..(50)  
<223> Xaa=Arg or Lys

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (88)..(88)

&lt;223&gt; Xaa=Val or Leu

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (109)..(109)

&lt;223&gt; Xaa=val or Leu

&lt;400&gt; 7

Xaa Val Val Met Thr Gln Xaa Pro Leu Xaa Leu Pro Val Thr Xaa Gly  
 1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Asp Ser  
 20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Leu Gln Gln Arg Pro Gly Gln Ser  
 35 40 45

Pro Xaa Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro  
 50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
 65 70 75 80

Ser Arg Val Glu Ala Glu Asp Xaa Gly Val Tyr Tyr Cys Trp Gln Gly  
 85 90 95

Thr His Phe Pro Arg Thr Phe Gly Gly Gly Thr Lys Xaa Glu Ile Lys  
 100 105 110

Arg

&lt;210&gt; 8

&lt;211&gt; 119

&lt;212&gt; PRT

&lt;213&gt; Artificial sequence

<220>

<223> humanized antibody

<220>

<221> MISC\_FEATURE

<222> (3)..(3)

<223> Xaa=Gln, Lys, or Arg

<220>

<221> MISC\_FEATURE

<222> (78)..(78)

<223> Xaa=Ser or Thr

<220>

<221> MISC\_FEATURE

<222> (87)..(87)

<223> Xaa=Arg or Lys

<220>

<221> MISC\_FEATURE

<222> (88)..(88)

<223> Xaa=Ala, Ser or Thr

<220>

<221> MISC\_FEATURE

<222> (114)..(114)

<223> Xaa=Leu, Thr, Ile, or Val

<400> 8

Glu Val Xaa Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Asn Tyr  
20 25 30



Gly Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
                   35                                  40                                  45

Ala Ser Ile Arg Ser Gly Gly Gly Arg Thr Tyr Tyr Ser Asp Asn Val  
           50                                  55                                  60

Lys Gly Arg Phe Thr Ile Ser Arg Glu Asn Ala Lys Asn Xaa Leu Tyr  
   65                                  70                                  75                                  80

Leu Gln Met Asn Ser Leu Xaa Xaa Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Val Arg Tyr Asp His Tyr Ser Gly Ser Ser Asp Tyr Trp Gly Gln Gly  
                   100                                  105                                  110

Thr Xaa Val Thr Val Ser Ser  
           115

<210> 9

<211> 113

<212> PRT

<213> Artificial sequence

<220>

<223> humanized antibody

<400> 9

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly  
   1                                  5                                  10                                  15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Asp Ser  
                   20                                  25                                  30

Asp Gly Lys Thr Tyr Leu Asn Trp Leu Gln Gln Arg Pro Gly Gln Ser  
           35                                  40                                  45

Pro Arg Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro  
           50                                  55                                  60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
   65                                  70                                  75                                  80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Trp Gln Gly  
                   85                                  90                                  95

Thr His Phe Pro Arg Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys  
 100 105 110

Arg

<210> 10

<211> 119

<212> PRT

<213> Artificial sequence

<220>

<223> humanized antibody

<400> 10

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Asn Tyr  
 20 25 30

Gly Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

Ala Ser Ile Arg Ser Gly Gly Gly Arg Thr Tyr Tyr Ser Asp Asn Val  
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Glu Asn Ala Lys Asn Ser Leu Tyr  
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

Val Arg Tyr Asp His Tyr Ser Gly Ser Ser Asp Tyr Trp Gly Gln Gly  
 100 105 110

Thr Leu Val Thr Val Ser Ser  
 115

<210> 11

<211> 219

<212> PRT

<213> Artificial sequence

<220>

<223> humanized antibody

<400> 11

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly  
1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Asp Ser  
20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Leu Gln Gln Arg Pro Gly Gln Ser  
35 40 45

Pro Arg Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro  
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Trp Gln Gly  
85 90 95

Thr His Phe Pro Arg Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys  
100 105 110

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu  
115 120 125

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe  
130 135 140

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln  
145 150 155 160

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser  
165 170 175

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu  
180 185 190

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser  
195 200 205

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
210 215

&lt;210&gt; 12

&lt;211&gt; 449

&lt;212&gt; PRT

&lt;213&gt; Artificial sequence

&lt;220&gt;

&lt;223&gt; humanized antibody

&lt;400&gt; 12

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Asn Tyr  
 20 25 30

Gly Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45

Ala Ser Ile Arg Ser Gly Gly Gly Arg Thr Tyr Tyr Ser Asp Asn Val  
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Glu Asn Ala Lys Asn Ser Leu Tyr  
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95

Val Arg Tyr Asp His Tyr Ser Gly Ser Ser Asp Tyr Trp Gly Gln Gly  
 100 105 110

Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe  
 115 120 125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu  
 130 135 140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp  
 145 150 155 160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu  
 165 170 175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser  
 180 185 190

Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro  
 195 200 205  
 Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys  
 210 215 220  
 Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro  
 225 230 235 240  
 Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser  
 245 250 255  
 Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp  
 260 265 270  
 Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn  
 275 280 285  
 Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val  
 290 295 300  
 Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu  
 305 310 315 320  
 Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys  
 325 330 335  
 Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr  
 340 345 350  
 Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr  
 355 360 365  
 Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu  
 370 375 380  
 Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu  
 385 390 395 400  
 Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys  
 405 410 415  
 Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu  
 420 425 430  
 Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly  
 435 440 445

Lys

&lt;210&gt; 13

&lt;211&gt; 46

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

&lt;223&gt; DNA primer

&lt;400&gt; 13

tatagagctc aagcttggat ggtggaaga tggatacagt tgggtgc

46

&lt;210&gt; 14

&lt;211&gt; 50

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

&lt;223&gt; DNA primer

&lt;400&gt; 14

tatagagctc aagcttccag tggatagach gatggggstg tygttttggc

50

&lt;210&gt; 15

&lt;211&gt; 1953

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

&lt;223&gt; humanized antibody

&lt;400&gt; 15

acgcgtccac catgatgagt cctgcccagt tcctgtttct gttagtgtc tggattcggg 60

aaaccaacgg tgatgttgtg atgaccagct ctccactctc cttgcctgtt accctgggac 120

aaccagcctc catctcttgc aagtcaagtc agagcctctt agatagtgat ggaaagacat 180

atttgaattg gttgcaacag cgcccaggcc agtctccaag acgcctaata tatctggtgt 240

```

ctaaactgga ctctggagtc cctgacaggt tctctggcag tggatcaggg acagatttta 300
cactgaaaat cagcagagtc gaggctgagg atgtgggagt ttattattgc tggcaaggta 360
cacatthttcc tcggacgttc ggtggaggca ccaagggtgga aatcaaactg aagtgcactt 420
tcctttctaga aattctaaac tctgaggggg tcggatgacg tggccattct ttgcctaaag 480
cattgagttt actgcaaggt cagaaaagca tgcaaagccc tcagaatggc tgcaaagagc 540
tccaacaaaa caatttagaa ctttattaag gaataggggg aagctaggaa gaaactcaaa 600
acatcaagat tttaaatacg cttcttggtc tccttgctat aattatctgg gataagcatg 660
ctgttttctg tctgtcccta acatgccctg tgattatccg caaacaacac acccaagggc 720
agaactttgt tacttaaaaca ccactctgtt tgcttctttc ctcaggaact gtggctgcac 780
catctgtctt catcttcccg ccactctgat agcagttgaa atctggaact gcctctgttg 840
tgtgcctgct gaataacttc tatcccagag aggccaaagt acagtggag gtggataacg 900
ccctccaatc gggtaactcc caggagagtg tcacagagca ggacagcaag gacagcacct 960
acagcctcag cagcaccctg acgctgagca aagcagacta cgagaaacac aaagtctacg 1020
cctgcgaagt caccatcag ggcctgagct cggccgtcac aaagagcttc aacaggggag 1080
agtgttagag ggagaagtgc cccacctgc tcctcagttc cagcctgacc ccctcccatc 1140
ctttggcctc tgaccctttt tccacagggg acctaccct attgcggtcc tccagctcat 1200
ctttcacctc acccccctcc tcctccttg ctttaattat gctaattgtg gaggagaatg 1260
aataaataaa gtgaatcttt gcacctgttg tttctctctt tcctcattta ataattatta 1320
tctgttggtt taccaactac tcaatttctc ttataaggga ctaaatatgt agtcaccta 1380
aggcgcataa ccatttataa aaatcatcct tcattctatt ttaccctatc atcctctgca 1440
agacagtcct ccctcaaacc cacaagcctt ctgtcctcac agtcccctgg gccatggtag 1500
gagagacttg cttccttggt tccccctcct cagcaagccc tcatagtcct ttttaagggt 1560
gacaggtctt acagtcatat atcctttgat tcaattcctt gagaatcaac caagcaaat 1620
ttttcaaaag aagaaacctg ctataaagag aatcattcat tgcaacatga tataaaataa 1680
caacacaata aaagcaatta aataaacaaa caatagggaa atgtttaagt tcatcatggt 1740
acttagactt aatggaatgt catgccttat ttacattttt aaacaggtac tgagggactc 1800
ctgtctgcca agggccgtat tgagtacttt ccacaacctt atttaatcca cactatactg 1860
tgagattaaa aacattcatt aaaatgttgc aaaggttcta taaagctgag agacaaatat 1920
attctataac tcagcaatcc cacttctagg atc 1953

```

&lt;210&gt; 16

&lt;211&gt; 3244

&lt;212&gt; DNA

&lt;213&gt; Artificial sequence

&lt;220&gt;

&lt;223&gt; humanized antibody

&lt;400&gt; 16

```

acgcgtccac catgaacttc gggctcagct tgattttcct tgccttgtc ttaaaagggtg      60
tccagtgtga agtgcaactg gtggagtctg ggggaggctt agtcagacct ggaggctctc      120
tgaggctctc ctgtgcaggc tctggattca ctttcagtaa ctatggcatg tcttggttctc      180
gccaggctcc tggaaaggga ctggagtggg ttgcatccat taggagtggg ggtggtagaa      240
cctactattc agacaatgta aagggccgat tcaccatctc cagagagaat gccaagaaca      300
gcctgtacct gcaaataaac agtctgagag ctgaggacac ggctgtctat tattgtgtca      360
gatatgatca ctatagtggg agctccgact actggggcca gggcaccttg gtcacagtct      420
cctcagggtga gtcctcaca cctctagagc tttctggggc aggccaggcc tgaccttggc      480
tttggggcag ggagggggct aaggtagggc aggtggcgcc agccagggtg acaccaatg      540
cccatgagcc cagacactgg acgctgaacc tcgcgacag ttaagaaccc aggggcctct      600
gcgccctggg ccagctctg tcccacaccg cggtcacatg gcaccacctc tcttgacgcc      660
tccaccaagg gcccatcggt cttccccctg gcacctcct ccaagagcac ctctgggggc      720
acagcggccc tgggctgcct ggtcaaggac tacttccccg aaccggtgac ggtgtcgtgg      780
aactcaggcg ccctgaccag cggcgtgcac accttccccg ctgtcctaca gtcctcagga      840
ctctactccc tcagcagcgt ggtgaccgtg ccctccagca gcttgggcac ccagacctac      900
atctgcaacg tgaatcaca gccagcaac accaagggtg acaagaaagt tggtagaggg      960
ccagcacagg gagggagggt gtctgtctga agccaggctc agcgctcctg cctggacgca     1020
tcccggctat gcagccccag tccagggcag caaggcaggc cccgtctgcc tcttcacccg     1080
gaggcctctg cccgccccac tcatgctcag ggagagggtc ttctggcttt tccccaggc     1140
tctgggcagg cacaggctag gtgcccctaa cccaggccct gcacacaaag gggcagggtg     1200
tgggctcaga cctgccaaga gccatatccg ggaggaccct gccctgacc taagcccacc     1260
ccaaaggcca aactctccac tccctcagct cggacacctt ctctcctccc agattccagt     1320
aactcccaat cttctctctg cagagcccaa atcttgtgac aaaactcaca catgcccacc     1380
gtgcccagggt aagccagccc aggcctcgcc ctccagctca aggcgggaca ggtgccctag     1440
agtagcctgc atccaggga aggcaccagc cgggtgctga cacgtccacc tccatctctt     1500
cctcagcacc tgaactcctg gggggaccgt cagtcttctt cttccccca aaaccaagg     1560
acaccctcat gatctcccg acccctgagg tcacatgcgt ggtggtggac gtgagccacg     1620

```



```

aagaccctga ggtcaagttc aactggtacg tggacggcgt ggaggtgcat aatgccaaaga 1680
caaagccgcg ggaggagcag tacaacagca cgtaccgtgt ggtcagcgtc ctcaccgtcc 1740
tgcaccagga ctggctgaat ggcaaggagt acaagtgcaa ggtctccaac aaagccctcc 1800
cagcccccat cgagaaaacc atctccaaag ccaaaggtag gacccgtggg gtgcgagggc 1860
cacatggaca gaggccggct cggcccaccc tctgccctga gagtgaccgc tgtaccaacc 1920
tctgtcccta cagggcagcc ccgagaacca caggtgtaca ccctgcccc atcccgggat 1980
gagctgacca agaaccagggt cagcctgacc tgcctggcca aaggcttcta tcccagcgac 2040
atcgccgtgg agtgggagag caatgggcag ccggagaaca actacaagac cagcctccc 2100
gtgctggact ccgacggctc cttcttcctc tacagcaagc tcaccgtgga caagagcagg 2160
tggcagcagg ggaacgtctt ctcatgtccc gtgatgcatg aggtctgca caaccactac 2220
acgcagaaga gcctctccct gtctccgggt aaatgagtgc gacggccggc aagccccgc 2280
tccccgggct ctgcgggtcg cagaggatg cttggcacgt acccctgta catacttccc 2340
gggcgccag catggaata aagcaccag cgctgccctg gggccctgcg agactgtgat 2400
ggttctttcc acgggtcagg ccgagtctga ggcctgagtg gcatgaggga ggcagagcgg 2460
gtcccactgt cccacactg gccaggctg tgcagggtg cctgggccgc ctagggtggg 2520
gctcagccag gggctgccct cggcaggggt ggggatttgc cagcgtggcc ctccctccag 2580
cagcacctgc cctgggctgg gccacgggaa gccctaggag cccctgggga cagacacaca 2640
ggccctgcct ctgtaggaga ctgtcctgtt ctgtgagcgc cctgtcctcc gacctccatg 2700
cccactcggg ggcattgcct gtccatgtgc gtagggacag gccctccctc acccatctac 2760
ccccacggca ctaaccctg gctgccctgc ccagcctgc acccgcattg ggacacaacc 2820
gactccgggg acatgcactc tcgggccctg tggagggaact ggtgcagatg cccacacaca 2880
cactcagccc agaccggtc aacaaacccc gactgaggt tggccggcca cagggccacc 2940
acacacacac gtgcacgcct cacacacgga gcctcaccg ggcgaactgc acagcaccca 3000
gaccagagca aggtcctcgc acacgtgaac actcctcggg cacaggcccc cagagcccc 3060
acgcggcacc tcaaggcca cgagcctctc ggcagcttct ccacatgtg acctgtcag 3120
acaaacccag cctcctctc acaagggtgc ccctgcagcc gccacacaca cacaggggat 3180
cacacaccac gtcacgtccc tggccctggc ccacttcca gtgccgccct tccctgcagg 3240
atcc 3244

```

&lt;210&gt; 17

&lt;211&gt; 717

&lt;212&gt; DNA

<213> Artificial sequence

<220>

<223> humanized antibody

<400> 17

```

atgatgagtc ctgcccagtt cctgtttctg ttagtgctct ggattcggga aaccaacggt      60
gatgttgtga tgaccagtc tccactctcc ttgcctgtta ccctgggaca accagcctcc      120
atctcttgca agtcaagtca gagcctctta gatagtgatg gaaagacata tttgaattgg      180
ttgcaacagc gcccaggcca gtctccaaga cgcctaattct atctggtgtc taaactggac      240
tctggagtc ctgacagggt ctctggcagt ggatcaggga cagattttac actgaaaatc      300
agcagagtcg aggctgagga tgtgggagtt tattattgct ggcaaggtag acattttcct      360
cggacgttcg gtggaggcac caaggaggaa atcaaactgt ctgtggctgc accatctgtc      420
ttcatcttcc cgccatctga tgagcagttg aaatctggaa ctgcctctgt tgtgtgcctg      480
ctgaataact tctatcccag agaggccaaa gtacagtggg aggtggataa cgccctcaa      540
tcgggtaact ccaggagag tgacacagag caggacagca aggacagcac ctacagcctc      600
agcagcacc cagcgtgag caaagcagac tacgagaac acaaagtcta cgcctgcgaa      660
gtcaccatc agggcctgag ctgcccgtc acaaagagct tcaacagggg agagtgt      717

```

<210> 18

<211> 1404

<212> DNA

<213> Artificial sequence

<220>

<223> humanized antibody

<400> 18

```

atgaacttcg ggctcagctt gattttcctt gtccttgtct taaaagggtg ccagtgtgaa      60
gtgcaactgg tggagtctgg gggaggctta gtgcagcctg gaggctctct gaggctctcc      120
tgtgcaggct ctggattcac tttcagtaac tatggcatgt cttgggttcg ccaggctcct      180
ggaaagggac tggagtgggt tgcattccatt aggagtgggt gtggtagaac ctactattca      240
gacaatgtaa agggccgatt caccatctcc agagagaatg ccaagaacag cctgtacctg      300
caaatgaaca gtctgagagc tgaggacacg gctgtctatt attgtgtcag atatgatcac      360
tatagtggta gctccgacta ctggggccag ggcaccttgg tcacagtctc ctacagcctc      420
accaagggcc catcgggtctt ccccctggca ccctcctcca agagcacctc tgggggcaca      480

```

gcggccctgg gctgcctggt caaggactac ttccccgaac cggtagcggg gtcgtggaac 540  
 tcaggcgccc tgaccagcgg cgtgcacacc ttcccgctg tcctacagtc ctcaggactc 600  
 tactccctca gcagcgtggt gaccgtgccc tccagcagct tgggcaccca gacctacatc 660  
 tgcaacgtga atcacaagcc cagcaacacc aaggtggaca agaaagttag gccccaatct 720  
 tgtgacaaaa ctcacacatg cccaccgtgc ccagcacctg aactcctggg gggaccgtca 780  
 gtcttcctct tcccccaaa acccaaggac accctcatga tctcccgac ccctgaggtc 840  
 acatgcgtgg tggtagcgtg gagccacgaa gaccctgagg tcaagttcaa ctggtacgtg 900  
 gacggcgtgg aggtgcataa tgccaagaca aagccgcggg agtagcagta caacagcacg 960  
 taccgtgtgg tcagcgtcct caccgtcctg caccaggact ggctgaatgg caaggagtac 1020  
 aagtgaagg tctccaacaa agccctccca gccccatcg agaaaaccat ctccaaagcc 1080  
 aaagggcagc cccgagaacc acaggtgtac accctgcccc catcccgga tgagctgacc 1140  
 aagaaccagg tcagcctgac ctgcctgggc aaaggcttct atcccagcga catcgccgtg 1200  
 gagtgggaga gcaatgggca gccggagaac aactacaaga ccacgcctcc cgtgctggac 1260  
 tccgacggct cttcttcct ctacagcaag ctcaccgtgg acaagagcag gtggcagcag 1320  
 gggaacgtct tctcatgctc cgtgatgcat gaggctctgc acaaccacta cacgcagaag 1380  
 agcctctccc tgtctccggg taaa 1404

<210> 19

<211> 239

<212> PRT

<213> Artificial sequence

<220>

<223> humanized antibody

<400> 19

Met Met Ser Pro Ala Gln Phe Leu Phe Leu Leu Val Leu Trp Ile Arg  
 1 5 10 15

Glu Thr Asn Gly Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro  
 20 25 30

Val Thr Leu Gly Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser  
 35 40 45

Leu Leu Asp Ser Asp Gly Lys Thr Tyr Leu Asn Trp Leu Gln Gln Arg  
 50 55 60

Pro Gly Gln Ser Pro Arg Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp  
65 70 75 80

Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe  
85 90 95

Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr  
100 105 110

Cys Trp Gln Gly Thr His Phe Pro Arg Thr Phe Gly Gly Gly Thr Lys  
115 120 125

Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro  
130 135 140

Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu  
145 150 155 160

Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp  
165 170 175

Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp  
180 185 190

Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys  
195 200 205

Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln  
210 215 220

Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
225 230 235

<210> 20

<211> 468

<212> PRT

<213> Artificial sequence

<220>

<223> humanized antibody

<400> 20

Met Asn Phe Gly Leu Ser Leu Ile Phe Leu Val Leu Val Leu Lys Gly  
1 5 10 15

Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln  
 20 25 30  
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe  
 35 40 45  
 Ser Asn Tyr Gly Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu  
 50 55 60  
 Glu Trp Val Ala Ser Ile Arg Ser Gly Gly Gly Arg Thr Tyr Tyr Ser  
 65 70 75 80  
 Asp Asn Val Lys Gly Arg Phe Thr Ile Ser Arg Glu Asn Ala Lys Asn  
 85 90 95  
 Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val  
 100 105 110  
 Tyr Tyr Cys Val Arg Tyr Asp His Tyr Ser Gly Ser Ser Asp Tyr Trp  
 115 120 125  
 Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro  
 130 135 140  
 Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr  
 145 150 155 160  
 Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr  
 165 170 175  
 Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro  
 180 185 190  
 Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr  
 195 200 205  
 Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn  
 210 215 220  
 His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser  
 225 230 235 240  
 Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu  
 245 250 255  
 Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu  
 260 265 270

Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser  
 275 280 285

His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu  
 290 295 300

Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr  
 305 310 315 320

Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn  
 325 330 335

Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro  
 340 345 350

Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln  
 355 360 365

Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val  
 370 375 380

Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val  
 385 390 395 400

Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro  
 405 410 415

Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr  
 420 425 430

Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val  
 435 440 445

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu  
 450 455 460

Ser Pro Gly Lys  
 465

```
*****
* User name:   mag (143)                               Queue:  LAH1/HP23-3      *
* File name:                               Server:  HP23-3      *
* Directory:                                         *
* Description: 557.pdf                               *
*               May 23, 2005                           10:43am    *
*****
*
*
*
*               mm m   aaa   ggg
*               m m m     a g   g
*               m m m   aaaa g   g
*               m   m a   a g   g
*               m   m   aaaa gggg
*                               g
*                               gggg
*
*****
*
*               L       SSS   TTTTT
*               L       S   S   T
*               L       S     T   ::
*               L       SSS   T   ::
*               L       S     T
*               L       S   S   T   ::
*               LLLLL   SSS   T   ::
*
*****
```

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 November 2002 (07.11.2002)

PCT

(10) International Publication Number  
**WO 02/088307 A2**

- (51) International Patent Classification<sup>7</sup>: C12N
- (21) International Application Number: PCT/US02/11854
- (22) International Filing Date: 26 April 2002 (26.04.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/287,653 30 April 2001 (30.04.2001) US
- (71) Applicant (for all designated States except US): **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **HINTON, Paul, Robert** [US/US]; 321 Dunsmuir Terrace, Number 4, Sunnyvale, CA 94086 (US). **VASQUEZ, Maximilano** [CR/US]; 3818 Louis Road, Palo Alto, CA 94303 (US).
- (74) Agents: **KELLEY, James, J. et al.**; **ELI LILLY AND COMPANY**, P. O. Box 6288, Indianapolis, IN 46206-6288 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/088307 A2

(54) Title: HUMANIZED ANTIBODIES

(57) Abstract: Humanized forms of mouse antibody 10D5 that retain the binding properties of mouse 10D5 are disclosed. Also disclosed are processes for making the humanized antibody, intermediates for making the humanized antibodies, including, nucleotide sequences, vectors, transformed host cells, and methods of using the humanized antibody to treat, prevent, alleviate, reverse, or otherwise ameliorate symptoms or pathology or both, that are associated with Down's syndrome or pre-clinical or clinical Alzheimer's disease or cerebral amyloid angiopathy.



## HUMANIZED ANTIBODIES

This application claims priority of US 60/287,653, filed 2001 April 30, the entire contents of which are incorporated herein by reference.

The invention relates to humanized antibodies useful for treating and preventing human diseases associated with amyloid  $\beta$  ( $A\beta$ ), such as Alzheimer's disease, Down's syndrome, and cerebral amyloid angiopathy. Mouse monoclonal antibody 10D5 was raised by immunizing mice with human  $A\beta$ 1-28, and has been widely used in analytical methods [*J. Neuropathol. Exper. Neurology* 51:76-83 (1992); *Nature* 359:325-327 (1992); *Neuroscience Lett.* 172:122-124 (1994); *Biochem. Biophys. Res. Commun.* 200:1598-1603 (1994); *J. Neuropathol. Exper. Neurology* 53:377-383 (1994); *Annals Neurology* 37:512-518 (1995); *Annals Neurology* 41:809-813 (1997); *J. Neuroimmunol.* 88:85-90 (1998); *J. Neuroimmunol.* 95:136-142 (1999)]. 10D5 has been shown to bind to the N-terminal region of  $A\beta$  and has affinity of approximately 43 pM for aggregated  $A\beta$ .

After 10D5 was administered to a group of 8.5 to 10.5 month-old heterozygous, transgenic PDAPP mice ( $APP^{V717F}$ ) at a weekly intraperitoneal dose of about 10 mg/kg for six months, the mice had significantly reduced levels of  $A\beta$ 1-42 in brain cortex. However, the 10D5 group did not have a significant reduction of total  $A\beta$  in any tissue, nor of  $A\beta$ 1-42 in hippocampus or cerebellum [Bard, F., *et al.*, *Nature Med.* 6:916-919 (2000); WO 00/72876 and WO 00/72880, 7 December, 2000]. It was asserted that amyloid plaques in the 10D5 group also reduced in number and appearance, with some evidence of cell-associated immunoreactivity.

Another study in WO 00/72876 and WO 00/72880 reported that administration of 10D5 to older mice for six months caused a significant reduction in amyloid  $\beta$  plaque burden. It was asserted that the antibody gained access to the central nervous system in sufficient amounts to "decorate"  $\beta$ -amyloid plaques. Finally, it was stated that mouse 10D5 induces phagocytosis of amyloid plaques in *in vitro* studies.

Methods for administering aggregated  $A\beta$ 1-42 to provoke an immunologic response and reduced amyloid deposits are described in PCT publication WO99/27944, published 10 June 1999. The description postulates that full-length aggregated  $A\beta$  peptide would be a useful immunogen. The application also indicates that antibodies that bind to  $A\beta$  peptide could be used as alternate therapeutic agents. However, this appears

to be speculation since the supporting data reflect protocols that involve active immunization using, for example, A $\beta$ 1-42.

WO 99/60024, published 25 November 1999, is directed to methods for amyloid removal using anti-amyloid antibodies. The mechanism, however, is stated to utilize the ability of anti-A $\beta$  antibodies to bind to pre-formed amyloid deposits (i.e. plaques) and result in subsequent microglial clearance of localized plaques. This mechanism was not proved *in vivo*. This publication further states that to be effective against A $\beta$  plaques, anti-A $\beta$  antibodies must be delivered directly to the brain, because antibodies cannot cross the blood brain barrier.

Queen, *et al.* describe methods of humanizing antibodies [*e.g.*, US Patent Nos. 5,585,089, 5,693,761, 5,693,762, 6,180,370].

Humanized forms of 10D5 are needed for use in humans having Down's syndrome, or pre-clinical or clinical Alzheimer's disease or cerebral amyloid angiopathy (CAA). However, it is not known whether 10D5 can be humanized so that the humanized antibody retained the binding properties of the mouse antibody.

#### Summary of the Invention

This invention provides humanized forms of 10D5. These humanized antibodies have binding properties (affinity and epitope location) that are approximately the same as those of the mouse 10D5 antibody. The invention includes antibodies, single chain antibodies, and fragments thereof. The invention includes antibodies wherein the CDR are those of mouse monoclonal antibody 10D5 (sequences SEQ ID NO:1 through SEQ ID NO:6) and wherein the antibodies retain approximately the binding properties of the mouse antibody and have *in vitro* and *in vivo* properties functionally equivalent to the mouse antibody. In another aspect, this invention provides humanized antibodies and fragments thereof, wherein the variable regions have sequences comprising the CDR from mouse antibody 10D5 and specific human framework sequences (sequences SEQ ID NO:7 - SEQ ID NO:10), wherein the antibodies retain approximately the binding properties of the mouse antibody and have *in vitro* and *in vivo* properties functionally equivalent to the mouse antibody 10D5. In another aspect, this invention provides humanized antibodies and fragments thereof, wherein the light chain is SEQ ID NO:11 and the heavy chain is SEQ ID NO:12.

Also part of the invention are polynucleotide sequences that encode the humanized antibodies or fragments thereof disclosed above, vectors comprising the polynucleotide sequences encoding the humanized antibodies or fragments thereof, host cells transformed with the vectors or incorporating the polynucleotides that express the humanized  
5 antibodies or fragments thereof, pharmaceutical formulations of the humanized antibodies and fragments thereof disclosed herein, and methods of making and using the same.

Such humanized antibodies and fragments thereof are useful for, among other things, treating and preventing diseases and conditions characterized by A $\beta$  plaques or A $\beta$  toxicity in the brain, such as Alzheimer's disease, Down's syndrome, and cerebral  
10 amyloid angiopathy in humans.

The invention also includes use of a humanized antibody of the present invention for the manufacture of a medicament, including prolonged expression of recombinant sequences of the antibody or antibody fragment in human tissues, for treating, preventing, or reversing Alzheimer's disease, Down's syndrome, or cerebral amyloid angiopathy, or  
15 to inhibit the formation of amyloid plaques or the effects of toxic soluble A $\beta$  species in humans.

#### Detailed Description of the Invention

We have surprisingly found that humanized antibodies, wherein the CDRs originate from mouse monoclonal antibody 10D5 and the framework and other portions  
20 of the antibodies originate from a human germ line, bind A $\beta$ 1-40 and A $\beta$ 1-42 with at least the affinity with which mouse 10D5 binds A $\beta$ . Thus, we have a reasonable basis for believing that humanized antibodies of this specificity, modified to reduce their immunogenicity by converting them to a humanized form, offer the opportunity to treat, both prophylactically and therapeutically, conditions in humans that are associated with  
25 formation of beta-amyloid plaques. These conditions include, as noted above, pre-clinical and clinical Alzheimer's, Down's syndrome, and pre-clinical and clinical cerebral amyloid angiopathy.

As used herein, the word "treat" includes therapeutic treatment, where a condition to be treated is already known to be present and prophylaxis - *i.e.*, prevention of, or  
30 amelioration of, the possible future onset of a condition.

By "antibody" is meant a monoclonal antibody *per se*, or an immunologically effective fragment thereof, such as an Fab, Fab', or F(ab')<sub>2</sub> fragment thereof. In some contexts, herein, fragments will be mentioned specifically for emphasis; nevertheless, it will be understood that regardless of whether fragments are specified, the term "antibody" includes such fragments as well as single-chain forms. As long as the protein retains the ability specifically to bind its intended target, it is included within the term "antibody." Also included within the definition "antibody" are single chain forms. Preferably, but not necessarily, the antibodies useful in the invention are produced recombinantly.

Antibodies may or may not be glycosylated, though glycosylated antibodies are preferred.

Antibodies are properly cross-linked via disulfide bonds, as is well known.

The basic antibody structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function.

Light chains are classified as kappa and lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, and define the antibody's isotype as IgG, IgM, IgA, IgD and IgE, respectively. Within light and heavy chains, the variable and constant regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 3 or more amino acids.

The variable regions of each light/heavy chain pair form the antibody binding site. Thus, an intact antibody has two binding sites. The chains all exhibit the same general structure of relatively conserved framework regions (FR) joined by three hypervariable regions, also called complementarity determining regions or CDRs. The CDRs from the two chains of each pair are aligned by the framework regions, enabling binding to a specific epitope. From N-terminal to C-terminal, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with well known conventions [Kabat, et al., "Sequences of Proteins of Immunological Interest" National Institutes of Health, Bethesda, Md., 1987 and 1991; Chothia, et al., J. Mol. Biol. 196:901-917 (1987); Chothia, et al., Nature 342:878-883 (1989)].

By "humanized antibody" is meant an antibody that is composed partially or fully of amino acid sequences derived from a human antibody germline by altering the sequence of an antibody having non-human complementarity determining regions (CDR). A humanized immunoglobulin does not encompass a chimeric antibody, having a mouse variable region and a human constant region. However, the variable region of the antibody and even the CDR are humanized by techniques that are by now well known in the art. The framework regions of the variable regions are substituted by the corresponding human framework regions leaving the non-human CDR substantially intact. As mentioned above, it is sufficient for use in the methods of the invention, to employ an immunologically specific fragment of the antibody, including fragments representing single chain forms.

Humanized antibodies have at least three potential advantages over non-human and chimeric antibodies for use in human therapy:

1) because the effector portion is human, it may interact better with the other parts of the human immune system (e.g., destroy the target cells more efficiently by complement-dependent cytotoxicity (CDC) or antibody-dependent cellular cytotoxicity (ADCC).

2) The human immune system should not recognize the framework or C region of the humanized antibody as foreign, and therefore the antibody response against such an injected antibody should be less than against a totally foreign non-human antibody or a partially foreign chimeric antibody.

3) Injected non-human antibodies have been reported to have a half-life in the human circulation much shorter than the half-life of human antibodies. Injected humanized antibodies will have a half-life essentially identical to naturally occurring human antibodies, allowing smaller and less frequent doses to be given.

The design of humanized immunoglobulins may be carried out as follows. As to the human framework region, a framework or variable region amino acid sequence of a CDR-providing non-human immunoglobulin is compared with corresponding sequences in a human immunoglobulin variable region sequence collection, and a sequence having a high percentage of identical amino acids is selected. When an amino acid falls under the following category, the framework amino acid of a human immunoglobulin to be used

(acceptor immunoglobulin) is replaced by a framework amino acid from a CDR-providing non-human immunoglobulin (donor immunoglobulin):

(a) the amino acid in the human framework region of the acceptor immunoglobulin is unusual for human immunoglobulin at that position, whereas the corresponding amino acid in the donor immunoglobulin is typical for human immunoglobulin at that position;

(b) the position of the amino acid is immediately adjacent to one of the CDRs; or

(c) any side chain atom of a framework amino acid is within about 5-6 angstroms (center-to-center) of any atom of a CDR amino acid in a three dimensional immunoglobulin model [Queen, *et al.*, Proc. Natl. Acad. Sci. USA 86:10029-10033 (1989), and Co, *et al.*, Proc. Natl. Acad. Sci. USA 88, 2869 (1991)]. When each of the amino acid in the human framework region of the acceptor immunoglobulin and a corresponding amino acid in the donor immunoglobulin is unusual for human immunoglobulin at that position, such an amino acid is replaced by an amino acid typical for human immunoglobulin at that position.

A preferred humanized antibody is a humanized form of mouse antibody 10D5.

The CDRs of humanized 10D5 have the following amino acid sequences:

light chain CDR1:

```

      1           5           10           15
20 Arg Ser Ser Gln Asn Ile Ile His Ser Asn Gly Asn Thr Tyr Leu Glu
   (SEQ ID NO:1)

```

light chain CDR2:

25      Lys Val Ser Asn Arg Phe Ser (SEQ ID NO:2)

light chain CDR3:

1 5  
Phe Gln Gly Ser His Val Pro Leu Thr (SEQ ID NO:3)

heavy chain CDR1:

1 5  
Thr Ser Gly Met Gly Val Ser (SEQ ID NO:4)

heavy chain CDR2:

His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser Leu Lys Ser  
(SEQ ID NO:5)

40 and, heavy chain CDR3:

Arg Pro Ile Thr Pro Val Leu Val Asp Ala Met Asp Tyr (SEQ ID NO:6).

A preferred light chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline Vk segment DPK18 and J segment Jk4:

```

5      1           5           10           15
   Asp Val Xaa Met Thr Gln Xaa Pro Leu Ser Leu Pro Val Xaa Leu Gly
      20           25           30
10    Xaa Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Asn Ile Xaa His Ser
      35           40           45
   Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
      50           55           60
15    Pro Xaa Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
      65           70           75           80
   Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
20    Ser Arg Val Glu Ala Glu Asp Xaa Gly Val Tyr Tyr Cys Phe Gln Gly
      85           90           95
   Ser His Val Pro Leu Thr Phe Gly Xaa Gly Thr Lys Xaa Glu Ile Lys
25
   Arg
                                     (SEQ ID NO:7) .

```

wherein:

- Xaa at position 3 is Val or Leu;
- 30 Xaa at position 7 is Ser or Thr;
- Xaa at position 14 is Thr or Ser;
- Xaa at position 17 is Gln, Asp, or Asn;
- Xaa at position 30 is Ile or Val;
- Xaa at position 50 is Arg or Lys;
- 35 Xaa at position 88 is Val or Leu;
- Xaa at position 105 is Gly or Ala; and
- Xaa at position 109 is Val or Leu.

A preferred heavy chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline VH segment DP-28 and J segment JH4, with several amino acid substitutions to the consensus amino acids in the same human subgroup to reduce potential immunogenicity:

```

1           5           10           15
Xaa Xaa Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu

           20           25           30
5  Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser

           35           40           45
   Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu

10          50           55           60
   Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Xaa

           65           70           75           80
   Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Xaa Xaa Gln Val

15          85           90           95
   Val Leu Xaa Xaa Thr Xaa Xaa Asp Pro Val Asp Thr Ala Thr Tyr Tyr

           100          105          110
20  Cys Val Arg Arg Pro Ile Thr Pro Val Leu Val Asp Ala Met Asp Tyr

           115          120
   Trp Gly Gln Gly Thr Xaa Val Thr Val Ser Ser

```

(SEQ ID NO:8).

25        wherein:

          Xaa at position 1 is Gln or Glu;  
          Xaa at position 2 is Val or Ala;  
          Xaa at position 64 is Ser or Thr;  
          Xaa at position 77 is Lys or Arg;  
30        Xaa at position 78 is Ser or Thr;  
          Xaa at position 83 is Thr or Ser;  
          Xaa at position 84 is Met, Ile, or Leu;  
          Xaa at position 86 is Asn, Ser, or Thr;  
          Xaa at position 87 is Met, Val, or Leu; and  
35        Xaa at position 118 is Leu or Ser.

A particularly preferred light chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline Vk segment DPK18 and J segment Jk4:

```

40  1           5           10           15
   Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly

           20           25           30
   Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Asn Ile Ile His Ser

45          35           40           45
   Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser

```



```

      50      55      60
Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro

5      65      70      75      80
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile

      85      90      95
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Phe Gln Gly

10
      100      105      110
Ser His Val Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys

15 Arg
                                     (SEQ ID NO:9)

```

A particularly preferred heavy chain variable region of a humanized antibody of the present invention has the following amino acid sequence, in which the framework originated from human germline VH segment DP-28 and J segment JH4:

```

1          5          10          15
Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu

25          20          25          30
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser

          35          40          45
Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu

30          50          55          60
Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser

          65          70          75          80
35 Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Ser Gln Val

          85          90          95
Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr

40          100          105          110
Cys Val Arg Arg Pro Ile Thr Pro Val Leu Val Asp Ala Met Asp Tyr

          115          120
Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser (SEQ ID NO:10).

45

```

A preferred light chain for a humanized antibody of the present invention has the  
50 amino acid sequence:

1 5 10 15  
Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly

20 25 30  
55 Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Asn Ile Ile His Ser

```

      35      40      45
Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser

      50      55      60
5  Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro

      65      70      75      80
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile

10  Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Phe Gln Gly

      100      105      110
15 Ser His Val Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys

      115      120      125
Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu

      130      135      140
20 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe

      145      150      155      160
Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln

25  Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser

      165      170      175
Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu

30  Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser

      180      185      190
      195      200      205
35  Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys (SEQ ID NO:11).
      210      215

```

A preferred heavy chain for a humanized antibody of the present invention has the amino acid sequence:

```

40  1      5      10      15
Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu

      20      25      30
45  Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser

      35      40      45
Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu

      50      55      60
50  Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser

      65      70      75      80
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Ser Gln Val

55  Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr

      85      90      95
Cys Val Arg Arg Pro Ile Thr Pro Val Leu Val Asp Ala Met Asp Tyr

60

```

115 120 125  
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly  
 130 135 140  
 5 Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly  
 145 150 155 160  
 Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val  
 10 165 170 175  
 Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe  
 180 185 190  
 Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val  
 15 195 200 205  
 Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val  
 210 215 220  
 20 Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys  
 225 230 235 240  
 Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu  
 25 245 250 255  
 Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr  
 260 265 270  
 Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val  
 30 275 280 285  
 Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val  
 290 295 300  
 35 Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser  
 305 310 315 320  
 Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu  
 40 325 330 335  
 Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala  
 340 345 350  
 Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro  
 45 355 360 365  
 Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln  
 370 375 380  
 50 Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala  
 385 390 395 400  
 Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr  
 55 405 410 415  
 Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu  
 420 425 430  
 Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser  
 60 435 440 445  
 Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser

450  
Leu Ser Pro Gly Lys

(SEQ ID NO:12) .

5

Other sequences are possible for the light and heavy chains for humanized 10D5. The immunoglobulins can have two pairs of light chain/heavy chain complexes, at least one chain comprising one or more mouse complementarity determining regions functionally joined to human framework region segments.

10

In another aspect, the present invention is directed to recombinant polynucleotides encoding antibodies which, when expressed, comprise the heavy and light chain CDRs from an antibody of the present invention. Exemplary polynucleotides, which on expression code for the polypeptide chains comprising the heavy and light chain CDRs of monoclonal antibody 10D5 are given herein. Due to codon degeneracy, other polynucleotide sequences can be readily substituted for those sequences. Particularly preferred polynucleotides of the present invention encode antibodies, which when expressed, comprise the CDRs of SEQ ID NO:1 – SEQ ID NO:6, or any of the variable regions of SEQ ID NO:7 – SEQ ID NO:10, or the light and heavy chains of SEQ ID NO:11 and SEQ ID NO:12.

15

20

The polynucleotides will typically further include an expression control polynucleotide sequence operably linked to the humanized immunoglobulin coding sequences, including naturally-associated or heterologous promoter regions. Preferably, the expression control sequences will be eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells, but control sequences for prokaryotic hosts may also be used. Once the vector has been incorporated into the appropriate host cell line, the host cell is propagated under conditions suitable for expressing the nucleotide sequences, and, as desired, the collection and purification of the light chains, heavy chains, light/heavy chain dimers or intact antibodies, binding fragments or other immunoglobulin forms may follow.

25

30

The nucleic acid sequences of the present invention capable of ultimately expressing the desired humanized antibodies can be formed from a variety of different polynucleotides (genomic or cDNA, RNA, synthetic oligonucleotides, etc.) and components (e.g., V, J, D, and C regions), using any of a variety of well known

techniques. Joining appropriate genomic and synthetic sequences is a common method of production, but cDNA sequences may also be utilized.

Below is a cDNA sequence (SEQ ID NO:17), from which the light chain having the amino acid sequence of SEQ ID NO:19 may be expressed.

```
5
  ATGAAGTTGCCTGTTAGGCTGTTGGTACTGATGTTCTGGATTCTGCTTCCAGCAGTGAT
  1 -----+-----+-----+-----+-----+-----+-----+ 60
    M K L P V R L L V L M F W I P A S S S D -
10
  GTTGTGATGACCCAATCTCCACTCTCCCTGCCTGTCACTCTTGGACAGCCAGCCTCCATC
  61 -----+-----+-----+-----+-----+-----+-----+ 120
    V V M T Q S P L S L P V T L G Q P A S I -
15
  TCTTGCAGATCTAGTCAGAACATTATACATAGTAATGGAAACACCTATTTAGAATGGTAC
  121 -----+-----+-----+-----+-----+-----+-----+ 180
    S C R S S Q N I I H S N G N T Y L E W Y -
20
  CTGCAGAAACCAGGCCAGTCTCCAAGGCTCCTGATCTACAAAGTTTCCAACCGATTTTCT
  181 -----+-----+-----+-----+-----+-----+-----+ 240
    L Q K P G Q S P R L L I Y K V S N R F S -
25
  GGGGTCCCAGACAGGTTCACTGGCAGTGGATCAGGGACAGATTTCACTCAAGATCAGC
  241 -----+-----+-----+-----+-----+-----+-----+ 300
    G V P D R F S G S G S G T D F T L K I S -
30
  AGAGTGGAGGCTGAGGATGTGGGAGTTTATTACTGCTTCAAGGTTACATGTTCCGCTC
  301 -----+-----+-----+-----+-----+-----+-----+ 360
    R V E A E D V G V Y Y C F Q G S H V P L -
35
  ACTTTCGGCGGAGGGACCAAGGTGAAATAAAACGAACTGTGGCTGCACCATCTGTCTTC
  361 -----+-----+-----+-----+-----+-----+-----+ 420
    T F G G G T K V E I K R T V A A P S V F -
40
  ATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGCTGCTG
  421 -----+-----+-----+-----+-----+-----+-----+ 480
    I F P P S D E Q L K S G T A S V V C L L -
45
  AATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAAGGTGGATAACGCCCTCCAATCG
  481 -----+-----+-----+-----+-----+-----+-----+ 540
    N N F Y P R E A K V Q W K V D N A L Q S -
50
  GGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGC
  541 -----+-----+-----+-----+-----+-----+-----+ 600
    G N S Q E S V T E Q D S K D S T Y S L S -
55
  AGCACCCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTC
  601 -----+-----+-----+-----+-----+-----+-----+ 660
    S T L T L S K A D Y E K H K V Y A C E V -
60
  ACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT (SEQ ID NO:17)
  661 -----+-----+-----+-----+-----+-----+-----+ 714
    T H Q G L S S P V T K S F N R G E C - (SEQ ID NO:19)
```

Below is a cDNA sequence (SEQ ID NO:18), from which the heavy chain having the amino acid sequence of SEQ ID NO:20 may be expressed.

```

  ATGGACAGGCTTACTTCCTCATTCTGCTGCTGATTGTCCCTGCATATGCTCTGTCCCAG
  1 -----+-----+-----+-----+-----+-----+-----+ 60
    M D R L T S S F L L L I V P A Y V L S Q -
```

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65

61 120  
V T L K E S G P V L V K P T E T L T L T -  
TGTACTTTCTCTGGGTTTTCACTGAGCACTTCTGGTATGGGAGTGAGCTGGATTCTGTCAG  
121 180  
C T F S G F S L S T S G M G V S W I R Q -  
CCTCCAGGAAAGGCTCTGGAGTGGCTGGCACACATTACTGGGATGATGACAAGCGCTAT  
181 240  
P P G K A L E W L A H I Y W D D D K R Y -  
AACCCATCCCTGAAGAGCCGGCTCACAATCTCCAAGGATACCTCCAAAAGCCAGGTAGTC  
241 300  
N P S L K S R L T I S K D T S K S Q V V -  
CTCACGATGACCAATATGGACCCTGTAGATACTGCCACATACTACTGTGTTCTGAAGGCCCC  
301 360  
L T M T N M D P V D T A T Y Y C V R R P -  
ATTACTCCGGTACTAGTCGATGCTATGGACTACTGGGGCCAAGGAACCCTGGTCACCGTC  
361 420  
I T P V L V D A M D Y W G Q G T L V T V -  
TCCTCAGCCTCCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCTCCTCCAAGAGCACC  
421 480  
S S A S T K G P S V F P L A P S S K S T -  
TCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACG  
481 540  
S G G T A A L G C L V K D Y F P E P V T -  
GTGTCTGGTGAACCTCAGGCGCCCTGACCAGCGGCGTGACACCTTCCCGGCTGTCTACAG  
541 600  
V S W N S G A L T S G V H T F P A V L Q -  
TCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACC  
601 660  
S S G L Y S L S S V V T V P S S S L G T -  
CAGACCTACATCTGCAACGTGAATCACAAGCCCAGCAACCAAGGTGGACAAGAAAGTT  
661 720  
Q T Y I C N V N H K P S N T K V D K K V -  
GAGCCCAAATCTGTGACAAAACCTCACACATGCCACCGTGCCCGACCACTGAACTCTTG  
721 780  
E P K S C D K T H T C P P C P A P E L L -  
GGGGGACCGTCAGTCTTCTCTTCCCCCAAACCCCAAGGACACCCTCATGATCTCCCGG  
781 840  
G G P S V F L F P P K P K D T L M I S R -  
ACCCCTGAGGTACATGCGTGGTGGTGACGTGAGCCACGAAGACCCTGAGGTCAAGTTC  
841 900  
T P E V T C V V V D V S H E D P E V K F -  
AACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAG  
901 960  
N W Y V D G V E V H N A K T K P R E E Q -  
TACAACAGCACGTACCGTGTGGTCAGCGTCCTACCGTCTGCACCAGGACTGGCTGAAT  
961 1020  
Y N S T Y R V V S V L T V L H Q D W L N -  
GGCAAGGAGTACAAGTGCAAGGTCTCCAACAAGCCCTCCAGCCCCCATCGAGAAAACC  
1021 1080  
G K E Y K C K V S N K A L P A P I E K T -

5  
 10  
 15  
 20  
 25

```

    ATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCTGCCCCATCCCGG
1081 -----+-----+-----+-----+-----+-----+-----+ 1140
    I S K A K G Q P R E P Q V Y T L P P S R -

    GATGAGCTGACCAAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGC
1141 -----+-----+-----+-----+-----+-----+-----+ 1200
    D E L T K N Q V S L T C L V K G F Y P S -

    GACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCACGCCT
1201 -----+-----+-----+-----+-----+-----+-----+ 1260
    D I A V E W E S N G Q P E N N Y K T T P -

    CCCGTGCTGGACTCCGACGGCTCCTTCTCTCTACAGCAAGCTCACCGTGGACAAGAGC
1261 -----+-----+-----+-----+-----+-----+-----+ 1320
    P V L D S D G S F F L Y S K L T V D K S -

    AGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCAC
1321 -----+-----+-----+-----+-----+-----+-----+ 1380
    R W Q Q G N V F S C S V M H E A L H N H -

    TACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA (SEQ ID NO:18)
1381 -----+-----+-----+-----+-----+-----+-----+ 1416
    Y T Q K S L S L S P G K - (SEQ ID NO:20)
  
```

The complete sequence of a humanized 10D5 light chain gene with introns  
 (located between MluI and BamHI sites, as in pVk-Hu10D5) is shown below (SEQ ID  
 NO:15). The nucleotide number indicates its position in pVk-Hu10D5. The V<sub>k</sub> and C<sub>k</sub>  
 exons are translated in single letter code; the dot indicates the translation termination  
 codon. The mature light chain starts at the double-underlined aspartic acid (D). The  
 intron sequences are in *italic*. The expressed light chain corresponds to SEQ ID NO:11  
 when mature.

35  
 40  
 45  
 50  
 55

```

    619 ACGCGTCCACCATGAAGTTGCCTGTTAGGCTGTTGGTACTGATGTTCTGGATTCCCTGCCAGCAGTGATGTTGTGATG
      M K L P V R L L V L M F W I P A S S S D V V M
    699 ACCCAATCTCCACTCTCCCTGCCTGTCTACTCTGGACAGCCAGCCTCCATCTCTTGCAGATCTAGTCAGAACATTATACA
      T Q S P L S L P V T L G Q P A S I S C R S S Q N I I H
    779 TAGTAATGGAACACCTATTTAGAAATGGTACCTGCAGAAACCAGGCCAGTCTCCAAGGCTCCTGATCTACAAAGTTTCCA
      S N G N T Y L E W Y L Q K P G Q S P R L L I Y K V S
    859 ACCGATTTTCTGGGGTCCCAGACAGGTTTCAGTGGCAGTGGAGTCAGGGACAGATTTACACTCAAGATCAGCAGAGTGGAG
      N R F S G V P D R F S G S G S G T D F T L K I S R V E
    939 GCTGAGGATGTGGGAGTTTATTACTGCTTTCAAGGTTTCACATGTTCCGCTCACTTTCGGCGGAGGGACCAAGGTGGAAAT
      A E D V G V Y Y C F Q G S H V P L T F G G G T K V E I
    1019 AAAACGTAAGTGCACTTTCTAATCTAGAAATCTAAACTCTGAGGGGGTGGATGACGTGGCCATTCTTTGCCTAAAGC
      K R
    1099 ATTGAGTTTACTGCAAGGTCAGAAAAGCATGCAAAGCCCTCAGAATGGCTGCAAAGAGCTCCAACAAAACAATTTAGAAC
    1179 TTTATTAAGGAATAGGGGGAAGCTAGGAAGAACTCAAACATCAAGATTTTAAATACGCTTCTTGGTCTCCTTGCTATA
    1259 ATTATCTGGGATAAGCATGCTGTTTCTGTCTGTCCCTAACATGCCCTGTGATTATCCGCAAAACAACACCCCAAGGGCA
    1339 GAACTTTGTACTTAAACACCATCTCTGTTTGCTTCTTTCCTCAGGAAGTGTGGCTGCACCATCTGTCTTTCATCTTCCCGC
      T V A A P S V F I F P
    1419 CATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCAGAGAGGCCAAAGTA
      P S D E Q L K S G T A S V V C L L N N F Y P R E A K V
    1499 CAGTGGAGGTTGGATAACGCCCTCAATCGGGTAAGTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTA
      Q W K V D N A L Q S G N S Q E S V T E Q D S K D S T Y
    1579 CAGCCTCAGCAGCACCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCAACCATCAGG
      S L S T L T L S K A D Y E K H K V Y A C E V T H Q
    1659 GCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTAGAGGGAGAAGTGCCCCACCTGCTCCTCAGTTCC
      G L S S P V T K S F N R G E C
    1739 AGCCTGACCCCTCCCATCTTTGGCCTCTGACCCCTTTTCCACAGGGGACCTACCCCTATTGCGGTCTCTCCAGCTCATC
    1819 TTTCACCTCACCCCTCTCTCTCTTGGCTTTAATTATGCTAATGTTGGAGGAGAATGAATAAA GTGAATCTTTTG
  
```

1899 CACCTGTGGTTTCTCTCTTCTCTCATTTAATAATTATATCTGTTGTTTACCAACTACTCAATTTCTCTTATAAGGGAC  
1979 TAAATATGTAGTCATCCTAAGGCGCATAACCAATTTATAAAAATCATCTTCATTTTACCCATCATCCTCTGCAA  
2059 GACAGTCCTCCCTCAAACCCACAAGCCTTCTGTCTCAGTCCCTGGGCCATGGTAGGAGAGACTTGCTTCTTGT  
2139 TCCCTCTCAGCAAGCCCTCATAGTCTTTTAAAGGGTGACAGGTCTTACAGTCATATATCCTTTGATTCAATTCCTG  
5 2219 AGAATCAACCAAAGCAAATTTTCAAAGAAGAAACCTGCTATAAAGAGAATCATTCATTGCAACATGATATAAAATAAC  
2299 AACACAATAAAAGCAATTAATAAACAACAATAGGGAAATGTTAAGTTCATCATGGTACTTAGACTTAATGGAATGTC  
2379 ATGCCTTATTACATTTTAAACAGGTACTGAGGACTCTGTCTGCCAAGGGCCGATTAGTACTTTCCACAACCTAA  
2459 TTTAATCCACACTATACTGTGAGATTAAAAACATTCATTAAAAATGTTGCAAGGTTCTATAAAGCTGAGAGACAAATATA  
2539 TTCTATAACTCAGCAATCCCCTTCTAGGATCC (SEQ ID NO:15)

10

The complete sequence of a humanized 10D5 heavy chain gene with introns  
(located between MluI and BamHI sites, as in pVg1-Hu10D5) is shown below (SEQ ID  
NO:16). The nucleotide number indicates its position in pVg1-Hu10D5. The V<sub>H</sub> and C<sub>H</sub>  
exons are translated in single letter code; the dot indicates the translation termination  
15 codon. The mature heavy chain starts at the double-underlined glutamine (Q). The intron  
sequences are in italic. The expressed heavy chain corresponds to SEQ ID NO:12 when  
mature.

619 ACGCGTCCACCATGGACAGGCTTACTTCCTCATTCTGCTGCTGATTGTCCCTGCATATGCTGTCAGGTTACTCTG  
20 M D R L T S S F L L L I V P A Y V L S Q = V T L  
699 AAAGAGTCTGGCCCTGTACTAGTGAAGCCACCGAGACCCTCACTCTGACTTGTACTTTCTCTGGGTTTTCACTGAGCAC  
K E S G P V L V K P T E T L T L T C T F S G F S L S T  
779 TTCTGGTATGGGAGTGAGCTGGATTCTGTCAGCCTCCAGGAAAGGCTCTGGAGTGGCTGGCACACATTTACTGGGATGATG  
S G M G V S W I R Q P P G K A L E W L A H I Y W D D  
25 859 ACAAGCGCTATAACCCATCCCTGAAGAGCCGGCTCACAATCTCCAAGGATACCTCCAAAGCCAGGTAGTCTCAGCATG  
D K R Y N P S L K S R L T I S K D T S K S Q V V L T M  
939 ACCAATATGGACCCTGTAGATACTGCCACATACTACTGTGTTGCAAGGCCCATTAATCCGGTACTAGTCGATGCTATGGA  
T N M D P V D T A T Y Y C V R R P I T P V L V D A M D  
1019 CTACTGGGGCCAAGGAACCCCTGGTCACCGTCTCCTCAGGTGAGTCTCACAACCTCTAGAGCTTTCTGGGGCAGGCCAGG  
30 Y W G Q G T L V T V S S  
1099 CCTGACCTTGGCTTTGGGGCAGGGAGGGGGCTAAGGTGAGGCGAGGTGGCGCCAGCCAGGTGCACACCCAATGCCCCATGAG  
1179 CCGAGACACTGGACGCTGAACCTCGCGGACAGTTAGGAACCCAGGGGCTCTGCGCCCTGGGCGCCAGCTCTGTCCACAC  
1259 CGCGGTACATGGCACCCACCTCTCTTGACGCTCCACCAAGGGCCCATCGTCTTCCCCCTGGCACCTCTCCAAGAGC  
A S T K G P S V F P L A P S S K S  
35 1339 ACCTCTGGGGGACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGTTGAAGTCAAG  
T S G G T A A L G C L V K D Y F P E P V T V S W N S G  
1419 CGCCCTGACCAGCGCGTGCACACCTTCCCGGCTGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCG  
A L T S G V H T F P A V L Q S S G L Y S L S S V V T  
1499 TGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAAGCCAGCAACCAAGGTGGACAGAGAA  
40 V P S S S L G T Q T Y I C N V N H K P S N T K V D K K  
1579 GTTGGTGAGAGGCCAGCACAGGGAGGGAGGTGTCTGCTGGAAGCCAGGCTCAGCGCTCTGCTGGACGCATCCCGGCT  
V  
1659 ATGACGCCCCAGTCCAGGGCAGCAAGGCAGGCCCCGTCTGCCTCTTCAACCGGAGGCTCTGCCCCCCCCACTCATGCTC  
1739 AGGGAGAGGCTCTTCTGGCTTTTCCCGAGGCTTGGGACGGCACAGGCTAGGTGCCCTTAACCCAGGCCCTGCACACAA  
45 1819 AGGGGAGGTGCTGGGCTCAGACCTGCCAAGAGCCATATCCGGGAGGACCTGCCCCTGACCTAAGCCCCACCCAAAGGC  
1899 CAAACTCTCCACTCCCTCAGCTCGGACACCTTCTCTCTCCAGATTCCAGTAACTCCCAATCTTCTCTGACAGGCC  
E P  
1979 AAATCTGTGACAAAACACACATGCCACCGTGCCAGGTAAGCCAGCCAGGCCCTCGCCCTCAGCTCAAGGCGGGA  
K S C D K T H T C P P C P  
50 2059 CAGGTGCCCTAGAGTAGCTGCATCCAGGGACAGGCCCCAGCCGGGTGCTGACACGTCCACCTCCATCTCTTCTCTCAGCA  
A  
2139 CCTGAACCTCTGGGGGACCGTCAGTCTTCTCTTCCCCCAAAACCCAAGGACACCCCTCATGATCTCCCGGACCCCTGA  
P E L L G G P S V F L F P P K P K D T L M I S R T P E  
2219 GGTACATGCGTGGTGGTGACGTGAGCCACGAAGACCTGAGGTCAAGTCAACTGGTACGTGGACGGCGTGGAGGTGC  
55 V T C V V V D V S H E D P E V K F N W Y V D G V E V  
2299 ATAATGCCAAGCAAAAGCCGCGGAGGAGCAGTACAACAGCAGTACCGTGTGGTCAGCGTCTCACCCTCTGCACCCAG  
H N A K T K P R E Q Y N S T Y R V V S V L H Q  
2379 GACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCAACAAAGCCCTCCAGCCCCATCGAGAAAACCATCTCCAA



D W L N G K E Y K C K V S N K A L P A P I E K T I S K  
 2459 AGCCAAAGGTGGGACCCGTGGGGTGGGAGGCCACATGGACAGAGGCCGGCTCGGCCACCCCTCTGCCCTGAGAGTGACC  
 A K  
 5 2539 GCTGTACCAACCTCTGTCCCTACAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCATCCCGGGATGAGCTGAC  
 G Q P R E P Q V Y T L P P S R D E L T  
 2619 CAAGAACCAGGTACAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGC  
 K N Q V S L T C L V K G F Y P S D I A V E W E S N G  
 2699 AGCCGGAGAACAACTACAAGACCACGCCTCCCGTGTGGACTCCGACGGCTCCTTCTCTCTACAGCAAGCTCACCGTG  
 Q P E N N Y K T T P P V L D S D G S F F L Y S K L T V  
 10 2779 GACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAA  
 D K S R W Q Q G N V P S C S V M H E A L H N H Y T Q K  
 2859 GAGCCTCTCCCTGTCTCCGGTAAATGAGTGCAGCGCCGGCAAGCCCCGCTCCCCGGGCTCTCGGGTCCGACGAGGA  
 S L S L S P G K -  
 2939 TGCTTGGCAGGTACCCCTGTACATACTTCCCGGGCGCCAGCATGGAAATAAA GCACCCAGCGCTGCCCTGGGCCCTG  
 15 3019 CGAGACTGTGATGGTCTTTCCACGGGTGAGGCCGAGTCTGAGGCCGTGAGTGGCATGAGGGAGGCAGAGCGGGTCCCACT  
 3099 GTCCCCACACTGGCCAGGCTGTGCAGGTGTGCCTGGGCCGCTAGGGTGGGGCTCAGCCAGGGGCTGCCCTCGGCAGGG  
 3179 TGGGGGATTGTCCAGCGTGGCCCTCCCTCCAGCAGCACCTGCCCTGGGCTGGGCCACGGGAAGCCCTAGGAGCCCCCTGGG  
 3259 GACAGACACACAGCCCTGCCTCTGTAGGAGACTGTCTGTCTGTGAGCGCCCTGTCTCCGACCTCCATGCCCACTCG  
 3339 GGGGCTGCCTAGTCCATGTGCGTAGGGACAGGCCCTCCCTCACCCTCTACCCCAAGGCACTAACCCCTGGCTGCCCT  
 20 3419 GCCCAGCCTCGCACCCGATGGGGACACAACCGACTCCGGGGACATGCACTCTCGGGCCCTGTGGAGGGACTGGTGCAGA  
 3499 TGCCACACACACACTCAGCCAGACCCGTTCAACAAACCCGCACTGAGGTTGGCCGGCCACAGGCCACACACACAC  
 3579 ACGTGCACGCCTCACACAGGAGCCTCACCCGGGCGAACTGCACAGCACCCAGACCAGAGCAAGGTCCTCGCACACGTGA  
 3659 ACACTCCTCGGACACAGGCCCCACGAGCCCCACGGGCACCTCAAGGCCACAGGCCTCTCGGCAGCTTCTCCACATGC  
 3739 TGACCTGCTCAGACAAACCCAGCCCTCTCTACAAGGGTGCCTTGCAGCCGCCACACACACAGGGGATCACACACC  
 25 3819 ACGTACAGTCCCTGGCCCTGGCCCACTTCCAGTGGCGCCCTTCCCTGCAGGATCC (SEQ ID NO:16)

Human constant region DNA sequences can be isolated in accordance with well  
 known procedures from a variety of human cells, but preferably from immortalized B-  
 cells. Suitable source cells for the polynucleotide sequences and host cells for  
 30 immunoglobulin expression and secretion can be obtained from a number of sources well-  
 known in the art.

In addition to the humanized immunoglobulins specifically described herein, other  
 "substantially homologous" modified immunoglobulins can be readily designed and  
 manufactured utilizing various recombinant DNA techniques well known to those skilled  
 35 in the art. For example, the framework regions can vary from the native sequences at the  
 primary structure level by several amino acid substitutions, terminal and intermediate  
 additions and deletions, and the like. Moreover, a variety of different human framework  
 regions may be used singly or in combination as a basis for the humanized  
 immunoglobulins of the present invention. In general, modifications of the genes may be  
 40 readily accomplished by a variety of well-known techniques, such as site-directed  
 mutagenesis.

Alternatively, polypeptide fragments comprising only a portion of the primary  
 antibody structure may be produced, which fragments possess one or more  
 immunoglobulin activities (e.g., complement fixation activity). These polypeptide  
 45 fragments may be produced by proteolytic cleavage of intact antibodies by methods well  
 known in the art, or by inserting stop codons at the desired locations in vectors using site-

directed mutagenesis, such as after CH1 to produce Fab fragments or after the hinge region to produce F(ab')<sub>2</sub> fragments. Single chain antibodies may be produced by joining VL and VH with a DNA linker.

As stated previously, the polynucleotides will be expressed in hosts after the sequences have been operably linked to (i.e., positioned to ensure the functioning of) an expression control sequence. These expression vectors are typically replicable in the host organisms either as episomes or as an integral part of the host chromosomal DNA. Commonly, expression vectors will contain selection markers, e.g., tetracycline or neomycin, to permit detection of those cells transformed with the desired DNA sequences.

*E. coli* is a prokaryotic host useful particularly for cloning the polynucleotides of the present invention. Other microbial hosts suitable for use include bacilli, such as *Bacillus subtilis*, and other enterobacteriaceae, such as *Salmonella*, *Serratia*, and various *Pseudomonas* species. In these prokaryotic hosts, one can also make expression vectors, which will typically contain expression control sequences compatible with the host cell (e.g., an origin of replication). In addition, any of a number of well-known promoters may be present, such as the lactose promoter system, a tryptophan (trp) promoter system, a beta-lactamase promoter system, or a promoter system from phage lambda. The promoters will typically control expression, optionally with an operator sequence, and have ribosome binding site sequences and the like, for initiating and completing transcription and translation.

Other microbes, such as yeast, may also be used for expression. *Saccharomyces* is a preferred host, with suitable vectors having expression control sequences, such as promoters, including 3-phosphoglycerate kinase or other glycolytic enzymes, and an origin of replication, termination sequences and the like as desired.

In addition to microorganisms, mammalian tissue cell culture may also be used to express and produce the polypeptides of the present invention. Eukaryotic cells are actually preferred, because a number of suitable host cell lines capable of secreting intact immunoglobulins have been developed in the art, and include the CHO cell lines, various COS cell lines, Syrian Hamster Ovary cell lines, HeLa cells, preferably myeloma cell lines, transformed B-cells, human embryonic kidney cell lines, or hybridomas. Expression vectors for these cells can include expression control sequences, such as an

origin of replication, a promoter, an enhancer, and necessary processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. Preferred expression control sequences are promoters derived from immunoglobulin genes, SV40, Adenovirus, Bovine Papilloma Virus, cytomegalovirus and the like.

The vectors containing the polynucleotide sequences of interest (e.g., the heavy and light chain encoding sequences and expression control sequences) can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment or electroporation may be used for other cellular hosts.

Once expressed, the antibodies can be purified according to standard procedures, including ammonium sulfate precipitation, ion exchange, affinity, reverse phase, hydrophobic interaction column chromatography, gel electrophoresis, and the like. Substantially pure immunoglobulins of at least about 90 to 95% homogeneity are preferred, and 98 to 99% or more homogeneity most preferred, for pharmaceutical uses. Once purified, partially or to homogeneity as desired, the polypeptides may then be used therapeutically or prophylactically, as directed herein.

The antibodies (including immunologically reactive fragments) are administered to a subject at risk for or exhibiting A $\beta$ -related symptoms or pathology such as clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical amyloid angiopathy, using standard administration techniques, preferably peripherally (*i.e.* not by administration into the central nervous system) by intravenous, intraperitoneal, subcutaneous, pulmonary, transdermal, intramuscular, intranasal, buccal, sublingual, or suppository administration. Although the antibodies may be administered directly into the ventricular system, spinal fluid, or brain parenchyma, and techniques for addressing these locations are well known in the art, it is not necessary to utilize these more difficult procedures. The antibodies of the invention are effective when administered by the more simple techniques that rely on the peripheral circulation system. The advantages of the present invention include the ability of the antibody to exert its beneficial effects even though not provided directly to the central nervous system itself. Indeed, it has been

demonstrated that the amount of antibody that crosses the blood-brain barrier is  $\leq 0.1\%$  of plasma levels.

The pharmaceutical compositions for administration are designed to be appropriate for the selected mode of administration, and pharmaceutically acceptable excipients such as, buffers, surfactants, preservatives, solubilizing agents, isotonicity agents, stabilizing agents and the like are used as appropriate. Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton PA, latest edition, incorporated herein by reference, provides a compendium of formulation techniques as are generally known to practitioners.

The concentration of the humanized antibody in formulations may range from as low as about 0.1% to as much as 15 or 20% by weight and will be selected primarily based on fluid volumes, viscosities, and so forth, in accordance with the particular mode of administration selected. Thus, a pharmaceutical composition for injection could be made up to contain in 1 mL of phosphate buffered saline from 1 to 100 mg of the humanized antibody of the present invention. The formulation could be sterile filtered after making the formulation, or otherwise made microbiologically acceptable. A typical composition for intravenous infusion could have a volume as much as 250 mL of fluid, such as sterile Ringer's solution, and 1-100 mg per mL, or more in antibody concentration. Therapeutic agents of the invention can be frozen or lyophilized for storage and reconstituted in a suitable sterile carrier prior to use. Lyophilization and reconstitution can lead to varying degrees of antibody activity loss (e.g. with conventional immune globulins, IgM antibodies tend to have greater activity loss than IgG antibodies). Dosages may have to be adjusted to compensate. The pH of the formulation will be selected to balance antibody stability (chemical and physical) and comfort to the patient when administered. Generally, pH between 4 and 8 is tolerated.

Although the foregoing methods appear the most convenient and most appropriate for administration of proteins such as humanized antibodies, by suitable adaptation, other techniques for administration, such as transdermal administration and oral administration may be employed provided proper formulation is designed.

In addition, it may be desirable to employ controlled release formulations using biodegradable films and matrices, or osmotic mini-pumps, or delivery systems based on dextran beads, alginate, or collagen.

In summary, formulations are available for administering the antibodies of the invention and are well-known in the art and may be chosen from a variety of options.

Typical dosage levels can be optimized using standard clinical techniques and will be dependent on the mode of administration and the condition of the patient.

5 The following examples are intended to illustrate but not to limit the invention.

The examples hereinbelow employ, among others, a murine monoclonal antibody designated "10D5" which was originally prepared by immunization with a peptide composed of residues 1-28 of human A $\beta$  peptide. As the examples here describe experiments conducted in murine systems, the use of murine monoclonal antibodies is satisfactory. However, in the treatment methods of the invention intended for human use, humanized forms of the antibodies with the immunospecificity corresponding to that of antibody 10D5 are preferred.

#### Example 1

##### Synthesis of Humanized Antibody 10D5

Cells and antibodies. Mouse myeloma cell line Sp2/0 was obtained from ATCC (Manassas, VA) and maintained in DME medium containing 10% FBS (Cat # SH30071.03, HyClone, Logan, UT) in a 37°C CO<sub>2</sub> incubator. Mouse 10D5 hybridoma cells were first grown in RPMI-1640 medium containing 10% FBS (HyClone), 10 mM HEPES, 2 mM glutamine, 0.1 mM non-essential amino acids, 1 mM sodium pyruvate, 25  $\mu$ g/ml gentamicin, and then expanded in serum-free media (Hybridoma SFM, Cat # 12045-076, Life Technologies, Rockville, MD) containing 2% low Ig FBS (Cat # 30151.03, HyClone) to a 1.5 liter volume in roller bottles. Mouse monoclonal antibody 10D5 (Mu10D5) was purified from the culture supernatant by affinity chromatography using a protein-G Sepharose column. Biotinylated Mu10D5 was prepared using EZ-Link Sulfo-NHS-LC-LC-Biotin (Cat # 21338ZZ, Pierce, Rockford, IL).

Cloning of variable region cDNAs. Total RNA was extracted from approximately 10<sup>7</sup> hybridoma cells using TRIzol reagent (Cat. # 15596-026, Life Technologies) and poly(A)<sup>+</sup> RNA was isolated with the PolyAtract mRNA Isolation System (Cat. # Z5310, Promega, Madison, WI) according to the suppliers' protocols. Double-stranded cDNA was synthesized using the SMART<sup>TM</sup>RACE cDNA Amplification Kit (Cat. # K1811-1, Clontech, Palo Alto, CA) following the supplier's protocol. The variable region cDNAs

for the light and heavy chains were amplified by polymerase chain reaction (PCR) using 3' primers that anneal respectively to the mouse kappa and gamma chain constant regions, and a 5' universal primer provided in the SMART<sup>TM</sup>RACE cDNA Amplification Kit. For VL PCR, the 3' primer has the sequence:

5

5' -TATAGAGCTCAAGCTTGGATGGTGGGAAGATGGATACAGTTGGTGC -3'

[SEQ ID NO:13]

with residues 17- 46 hybridizing to the mouse Ck region. For VH PCR, the 3' primers  
10 have the degenerate sequences:

5' -TATAGAGCTCAAGCTTCCAGTGGATAGACCGATGGGGCTGTCGTTTTGGC -3'   
A G T  
T

15 [SEQ ID NO:14]

with residues 17 - 50 hybridizing to mouse gamma chain CH1. The VL and VH cDNAs were subcloned into pCR4Blunt-TOPO vector (Cat. # 45-0031, Invitrogen, Carlsbad, CA) for sequence determination. DNA sequencing was carried out by PCR cycle sequencing  
20 reactions with fluorescent dideoxy chain terminators (Applied Biosystems, Foster City, CA) according to the manufacturer's instructions. The sequencing reactions were analyzed on a Model 377 DNA Sequencer (Applied Biosystems).

Construction of humanized 10D5 (Hu10D5) variable regions. Humanization of the mouse antibody V regions was carried out as outlined by Queen et al., 1989, *op. Cit.* The human V region framework used as acceptor for Mu10D5 CDRs was chosen based on sequence homology. The computer programs ABMOD and ENCAD [Levitt, M., J. Mol. Biol. 168:595-620 (1983)] were used to construct a molecular model of the variable regions. Amino acids in the humanized V regions that were predicted to have contact with CDRs were substituted with the corresponding residues of Mu10D5. This was done at residue 98 in the heavy chain and at residues 41 and 51 in the light chain. The amino acids in the humanized V region that were found to be rare in the same V-region subgroup were changed to the consensus amino acids to eliminate potential immunogenicity. This was done at residues 42 and 44 in the light chain and at residue 24 in the heavy chain.

The light and heavy chain variable region genes were constructed and amplified using eight overlapping synthetic oligonucleotides ranging in length from approximately 65 to 80 bases [He, X. Y., et al., J. Immunol. 160: 1029-1035 (1998)]. The oligonucleotides were annealed pairwise and extended with the Klenow fragment of DNA polymerase I, yielding four double-stranded fragments. The resulting fragments were denatured, annealed pairwise, and extended with Klenow, yielding two fragments. These fragments were denatured, annealed pairwise, and extended once again, yielding a full-length gene. The resulting product was amplified by PCR using the Expand High Fidelity PCR System (Cat. # 1 732 650, Roche Molecular Biochemicals, Indianapolis, IN). The PCR-amplified fragments were gel-purified and cloned into pCR4Blunt-TOPO vector. After sequence confirmation, the VL and VH genes were digested with MluI and XbaI, gel-purified, and subcloned respectively into vectors for expression of light and heavy chains to make pVk-Hu10D5 and pVg1-Hu10D5 [Co, M. S., et al., J. Immunol. 148:1149-1154 (1992)]. The mature humanized 10D5 antibody expressed from these plasmids has the light chain of SEQ ID NO:11 and the heavy chain of SEQ ID NO:12.

Stable transfection. Stable transfection into mouse myeloma cell line Sp2/0 was accomplished by electroporation using a Gene Pulser apparatus (BioRad, Hercules, CA) at 360 V and 25  $\mu$ F as described (Co, et al., 1992, *op. cit.*). Before transfection, pVk-Hu10D5 and pVg1-Hu10D5 plasmid DNAs were linearized using FspI. Approximately  $10^7$  Sp2/0 cells were transfected with 20  $\mu$ g of pVk-Hu10D5 and 40  $\mu$ g of pVg1-Hu10D5. The transfected cells were suspended in DME medium containing 10% FBS and plated

into several 96-well plates. After 48 hr, selection media (DME medium containing 10% FBS, HT media supplement, 0.3 mg/ml xanthine and 1 µg/ml mycophenolic acid) was applied. Approximately 10 days after the initiation of the selection, culture supernatants were assayed for antibody production by ELISA as shown below. High yielding clones were expanded in DME medium containing 10% FBS and further analyzed for antibody expression. Selected clones were then adapted to growth in Hybridoma SFM.

Measurement of antibody expression by ELISA. Wells of a 96-well ELISA plate (Nunc-Immuno plate, Cat # 439454, NalgeNunc, Naperville, IL) were coated with 100 µl of 1 µg/ml goat anti-human IgG, Fc γ fragment specific, polyclonal antibodies (Cat. # 109-005-098, Jackson ImmunoResearch, West Grove, PA) in 0.2 M sodium carbonate-bicarbonate buffer (pH 9.4) overnight at 4°C. After washing with Washing Buffer (PBS containing 0.1% Tween 20), wells were blocked with 400 µl of Superblock Blocking Buffer (Cat # 37535, Pierce) for 30 min and then washed with Washing Buffer. Samples containing Hu10D5 were appropriately diluted in ELISA Buffer (PBS containing 1% BSA and 0.1% Tween 20) and applied to ELISA plates (100 µl per well). As a standard, humanized anti-CD33 IgG1 monoclonal antibody HuM195 (Co, *et al.*, 1992, *op. cit.*) was used. The ELISA plate was incubated for 2 hr at room temperature and the wells were washed with Washing Buffer. Then, 100 µl of 1/1,000-diluted HRP-conjugated goat anti-human kappa polyclonal antibodies (Cat # 1050-05, Southern Biotechnology, Birmingham, AL) in ELISA Buffer was applied to each well. After incubating for 1 hr at room temperature and washing with Washing Buffer, 100 µl of ABTS substrate (Cat #s 507602 and 506502, Kirkegaard and Perry Laboratories, Gaithersburg, MD) was added to each well. Color development was stopped by adding 100 µl of 2% oxalic acid per well. Absorbance was read at 415 nm using an OPTImax microplate reader (Molecular Devices, Menlo Park, CA).

Purification of Hu10D5. One of the high Hu10D5-expressing Sp2/0 stable transfectants (clone #1) was adapted to growth in Hybridoma SFM and expanded to 2 liters in roller bottles. Spent culture supernatant was harvested when cell viability reached 10% or below and loaded onto a protein-A Sepharose column. The column was washed with PBS before the antibody was eluted with 0.1 M glycine-HCl (pH 2.8), 0.1 M NaCl. The eluted protein was dialyzed against 3 changes of 2 liters of PBS and filtered through a 0.2 µm filter prior to storage at 4°C. Antibody concentration was determined



by measuring absorbance at 280 nm ( $1 \text{ mg/ml} = 1.4 A_{280}$ ). SDS-PAGE in Tris-glycine buffer was performed according to standard procedures on a 4-20% gradient gel (Cat # EC6025, Novex, San Diego, CA). Purified humanized 10D5 antibody is reduced and run on an SDS- PAGE gel. The whole antibody shows two bands of approximate molecular weights 25 kDa and 50 kDa. These results are consistent with the molecular weights of the light chain and heavy chain, or with the molecular weight of the chain(s) comprising a fragment, calculated from their amino acid compositions.

### Example 2

#### *In vitro* binding properties of humanized 10D5 antibody

The binding efficacy of humanized 10D5 antibody, synthesized and purified as described above, was compared with the mouse 10D5 antibody using biotinylated mouse 10D5 antibody in a comparative ELISA. Wells of a 96-well ELISA plate (Nunc-Immuno plate, Cat # 439454, NalgeNunc) were coated with 100  $\mu\text{l}$  of  $\beta$ -amyloid peptide (1-42) in 0.2 M sodium carbonate/bicarbonate buffer (pH 9.4) (1  $\mu\text{g/mL}$ ) overnight at 4°C.

After washing the wells with phosphate buffered saline (PBS) containing 0.1% Tween 20 (Washing Buffer) using an ELISA plate washer, the wells were blocked by adding 300  $\mu\text{L}$  of SuperBlock reagent (Pierce) per well. After 30 minutes of blocking, the wells were washed with Washing Buffer and excess liquid was removed.

A mixture of biotinylated Mu10D5 (0.4  $\mu\text{g/ml}$  final concentration) and competitor antibody (Mu10D5 or Hu10D5; starting at 1000  $\mu\text{g/ml}$  final concentration and serial 3-fold dilutions) in ELISA Buffer were added in triplicate in a final volume of 100  $\mu\text{l}$  per well. As a no-competitor control, 100  $\mu\text{l}$  of 0.4  $\mu\text{g/ml}$  biotinylated Mu10D5 was added. As a background control, 100  $\mu\text{l}$  of ELISA Buffer was added. The ELISA plate was incubated at room temperature for 90 min. After washing the wells with Washing Buffer, 100  $\mu\text{l}$  of 10  $\mu\text{g/ml}$  HRP-conjugated streptavidin (Cat # 21124, Pierce) was added to each well. The plate was incubated at room temperature for 30 min and washed with Washing Buffer. For color development, 100  $\mu\text{l/well}$  of ABTS Peroxidase Substrate (Kirkegaard & Perry Laboratories) was added. Color development was stopped by adding 100  $\mu\text{l/well}$  of 2% oxalic acid. Absorbance was read at 415 nm. The absorbances were plotted against the log of the competitor concentration, curves were fit to the data points (using Prism) and the IC50 was determined for each antibody using methods well-known in the art.

The mean IC<sub>50</sub> for mouse 10D5 was 23.4 µg/mL (three separate experiments, standard deviation = 5.5 µg/mL) and for humanized 10D5 was 49.1 µg/mL (three separate experiments, standard deviation = 11.8 µg/mL). A second set of three experiments was carried out, essentially as described above, and the mean IC<sub>50</sub> for mouse 10D5 was  
5 determined to be 20 µg/mL (SD = 1 µg/mL) and for humanized 10D5, the IC<sub>50</sub> was determined to be 16 µg/mL (SD = 0.6 µg/mL). On the basis of these results, we conclude that humanized 10D5 has binding properties that are very similar to those of the mouse antibody 10D5. Therefore, we expect that humanized 10D5 has very similar *in vitro* and *in vivo* activities compared with mouse 10D5 and will exhibit in humans the same effects  
10 demonstrated with mouse 10D5 in mice.

### Example 3

#### *In vitro* binding properties of mouse and humanized antibodies 10D5

Antibody affinity ( $KD = K_d / K_a$ ) was determined using a BIAcore biosensor 2000  
15 and data analyzed with BIAevaluation (v. 3.1) software. A capture antibody (rabbit anti-mouse Ig or anti-human Ig) was coupled via free amine groups to carboxyl groups on flow cell 2 of a biosensor chip (CM5) using N-ethyl-N-dimethylaminopropyl carbodiimide and N-hydroxysuccinimide (EDC/NHS). A non-specific rabbit IgG was coupled to flow cell 1 as a background control. Monoclonal antibodies were captured to yield 300 resonance  
20 units (RU). Amyloid-beta 1-40 or 1-42 (Biosource International, Inc.) was then flowed over the chip at decreasing concentrations (1000 to 0.1 times KD). To regenerate the chip, bound anti-A $\beta$  antibody was eluted from the chip using a wash with glycine-HCl (pH 2). A control injection containing no amyloid-beta served as a control for baseline subtraction. Sensorgrams demonstrating association and dissociation phases were  
25 analyzed to determine  $K_d$  and  $K_a$ . The affinity (KD) of mouse antibody 10D5 for A $\beta$  1-40 was determined to be 390 nM, and the affinity of humanized 10D5, prepared essentially as described in Example 1, was determined to be 209 nM. Affinity for A $\beta$  1-42 was biphasic for both mouse 10D5 and humanized 10D5. For mouse 10D5, the affinities for A $\beta$  1-42 were 0.57 nM and 4950 nM. Humanized 10D5 had affinities for  
30 A $\beta$  1-42 of 0.19 nM and 1020 nM.

#### Example 4

##### Epitope mapping of mouse and humanized 10D5

The BIAcore is an automated biosensor system for measuring molecular interactions [Karlsson R., *et al. J. Immunol. Methods* 145:229-240 (1991)]. The advantage of the BIAcore over other binding assays is that binding of the antigen can be measured without having to label or immobilize the antigen (i.e. the antigen maintains a more native conformation). The BIAcore methodology was used to assess the binding of various amyloid-beta peptide fragments to either mouse 10D5 or humanized 10D5 (prepared substantially as described in Example 1). All dilutions were made with HEPES buffered saline containing Tween 20. A single concentration of a variety of fragments of human A $\beta$  or mouse A $\beta$  1-40 (BioSource International) was used. Human amyloid beta fragments 1-10 and 1-20 bound to mouse 10D5 and to humanized 10D5, while human A $\beta$  fragments 10-20 and 16-25 did not bind to either antibody. Neither mouse 10D5 nor humanized 10D5 bound mouse A $\beta$  1-40. Using this methodology, the binding epitope for both mouse and humanized 10D5 appears to be between amino acids 1 and 10 of human A $\beta$ .

#### Example 5

##### In vivo experiments with 10D5

Unless otherwise stated, all studies used PDAPP mice, and all injections were intraperitoneal (i.p.) In general, a control group of mice received injections of saline. In some cases, another control group received injections of a non-specific mouse IgG preparation.

Six weeks of weekly injection of 360  $\mu$ g of 10D5 in old mice (24 month) raised soluble A $\beta_{\text{total}}$  in hippocampus by 16% and A $\beta$  1-42 in hippocampus by 21%, while lowering hippocampal insoluble A $\beta_{\text{total}}$  by 24% and A $\beta$  1-42 by 26% (no statistically significant difference; 9 animals per control group and 10 animals per antibody group). In the cortex, mean insoluble A $\beta_{\text{total}}$  was lower by 27% and A $\beta$  1-42 by 29%, while mean insoluble A $\beta$  1-40 increased by 7% (no statistically significant differences).

In hemizygous, 4 month old mice, administration of 360  $\mu$ g 10D5 per animal: 1) raised average plasma A $\beta$  1-40 and A $\beta$  1-42 levels approximately 3-fold by 24 hours after administration; and 2) had no significant effect on soluble A $\beta$  1-40 in the cortex after 24

hours compared with saline control (no differences were statistically significant; 5 animals per group).

Administration of 360  $\mu$ g of 10D5 per animal (5 animals per group, saline control): 1) raised average plasma A $\beta$  1-40 and A $\beta$  1-42 levels approximately 14-fold and 19-fold, respectively by 24 hours after administration; 2) had no consistent or significant effect on soluble or insoluble A $\beta$  1-40, A $\beta$  1-42, or A $\beta_{\text{total}}$  in the cortex or hippocampus after 24 hours; 3) lowered soluble A $\beta$  1-40, A $\beta$  1-42, and A $\beta_{\text{total}}$  in the cerebellum by 50% ( $p < 0.05$ ), 33%, and 13%, respectively; and 4) lowered insoluble A $\beta$  1-40, A $\beta$  1-42, and A $\beta_{\text{total}}$  in the cerebellum by 53% ( $p < 0.001$ ), 46% ( $p < 0.001$ ), and 30% ( $p < 0.01$ ), respectively.

In young mice, administration of 360  $\mu$ g of 10D5 per animal (5 per group): 1) raised average plasma A $\beta$  1-42 levels approximately 33% by 24 hours after administration; and 2) in the cortex, raised soluble A $\beta$  1-40 3.4-fold ( $p < 0.001$ ), lowered soluble A $\beta$  1-42 by 22% ( $p < 0.05$ ), lowered insoluble A $\beta$  1-40 about 10% and increased insoluble A $\beta$  1-42 about 12%.

We claim:

1. Humanized 10D5 antibody.

2. A humanized antibody, or fragment thereof, comprising a humanized light chain comprising three light chain complementarity determining regions (CDRs) from the mouse monoclonal antibody 10D5 and a light chain variable region framework sequence from a human immunoglobulin light chain; and a humanized heavy chain comprising three heavy chain CDRs from the mouse monoclonal antibody 10D5 and a heavy chain variable region framework sequence from a human immunoglobulin heavy chain; wherein the light chain CDRs have the following amino acid sequences:

10 light chain CDR1:

1 5 10 15  
Arg Ser Ser Gln Asn Ile Ile His Ser Asn Gly Asn Thr Tyr Leu Glu  
(SEQ ID NO:1)

15 light chain CDR2:

1 5  
Lys Val Ser Asn Arg Phe Ser (SEQ ID NO:2)

light chain CDR3:

20 1 5  
Phe Gln Gly Ser His Val Pro Leu Thr (SEQ ID NO:3)

and the heavy chain CDRs have the following amino acid sequences:

heavy chain CDR1:

25 1 5  
Thr Ser Gly Met Gly Val Ser (SEQ ID NO:4)

heavy chain CDR2:

30 1 5 10 15  
His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser Leu Lys Ser  
(SEQ ID NO:5)

and, heavy chain CDR3:

35 1 5 10  
Arg Pro Ile Thr Pro Val Leu Val Asp Ala Met Asp Tyr (SEQ ID NO:6).

3. A humanized antibody or fragment thereof comprising a humanized light chain variable region having the sequence of SEQ ID NO:7 and a humanized heavy variable region having the sequence of SEQ ID NO:8.

4. The humanized antibody or fragment thereof of claim 3 having a light chain variable region of the sequence given by SEQ ID NO:9 and a heavy chain variable region given by SEQ ID NO:10.

5. The humanized antibody or fragment thereof of claim 3 having a light chain of the sequence given by SEQ ID NO:11 and a heavy chain of the sequence given by SEQ ID NO:12.

6. An antibody fragment obtainable by enzymatic cleavage of the humanized antibody of any one of claims 1 - 5.

7. An Fab or F(ab')<sub>2</sub> fragment of any one of the humanized antibodies of claims 1 - 5.

8. The F(ab')<sub>2</sub> fragment of claim 7.

9. The Fab fragment of claim 7.

10. The humanized antibody or fragment of any one of claims 1 - 9, which is a single chain antibody.

11. The humanized antibody or fragment of any one of claims 1 - 10 that is an IgG<sub>1</sub> immunoglobulin isotype.

12. The humanized antibody or fragment of any one of claims 1 - 11, wherein the antibody or fragment thereof is produced in a host cell selected from the group consisting of a myeloma cell, a chinese hamster ovary cell, a syrian hamster ovary cell, and a human embryonic kidney cell.

13. A polynucleotide compound, comprising a sequence coding for the light chain or the heavy chain of the humanized antibody of any one of claims 1 - 12, or a fragment thereof.

14. A polynucleotide sequence, which when expressed in a suitable host cell, yields an antibody of any one of claims 1 – 12.

15. The polynucleotide of claim 13 or 14 selected from the group consisting of SEQ ID NO: 15, SEQ ID NO: 17, and a polynucleotide comprising a sequence that codes  
5 for the light chain variable region given by SEQ ID NO:7, SEQ ID NO:9, or SEQ ID NO: 11.

16. The polynucleotide of claim 13 or 14 selected from the group consisting of SEQ ID NO:16, SEQ ID NO:18, and a polynucleotide comprising a sequence that codes for the heavy chain variable region given by SEQ ID NO:8, SEQ ID NO:10, or SEQ ID  
10 NO:12.

17. An expression vector for expressing the antibody of any one of claims 1 - 12 comprising the polynucleotide sequence of any one of claims 13 - 16.

18. A cell transfected with the expression vector of claim 17.

19. A cell transfected with two expression vectors of claim 17, wherein a first  
15 vector comprises the polynucleotide sequence coding for the light chain and a second vector comprises the sequence coding for the heavy chain.

20. A cell that is capable of expressing the humanized antibody or fragment of any one of claims 1 – 12.

21. The cell of any one of claims 18 – 20, wherein the cell is selected from the  
20 group consisting of a myeloma cell, a chinese hamster ovary cell, a syrian hamster ovary cell, and a human embryonic kidney cell.

22. A pharmaceutical composition comprising the humanized antibody or fragment of any one of claims 1 – 12, and a pharmaceutically acceptable excipient.

23. A method of treating Down's syndrome, clinical or pre-clinical Alzheimer's disease, or clinical or pre-clinical cerebral amyloid angiopathy in a human subject, comprising administering to the human subject an effective amount of a humanized antibody or fragment of any one of claims 1 – 12.

5        24. A method to inhibit the formation of A $\beta$  plaque in the brain of a human subject, comprising administering to the human subject an effective amount of the humanized antibody or fragment of any one of claims 1 – 12.

25. A method to reduce A $\beta$  plaque in the brain of a human subject, comprising administering to the human subject an effective amount of a humanized antibody or  
10       fragment of any one of claims 1 – 12.

26. The method of either of claims 24 – 25, wherein the subject is diagnosed with clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical cerebral amyloid angiopathy.

27. The method of any one of claims 24 – 25, wherein the subject is not  
15       diagnosed with clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical cerebral amyloid angiopathy.

28. Use of the humanized antibody or a fragment thereof according to any one of Claims 1 – 12 for the manufacture of a medicament, including prolonged expression of recombinant sequences of the antibody or antibody fragment in human tissues, for treating  
20       clinical or pre-clinical Alzheimer's disease, Down's syndrome, or clinical or pre-clinical cerebral amyloid angiopathy.

29. Use of the humanized antibody or fragment of any one of claims 1 – 12 for the manufacture of a medicament for treating Alzheimer's disease.



## SEQUENCE LISTING

<110> Eli Lilly and Company

<120> Humanized Antibodies

<130> X-14819

<150> US 60/287,653

<151> 2001-04-30

<160> 20

<170> PatentIn version 3.1.

<210> 1

<211> 16

<212> PRT

<213> mus sp.

<400> 1

Arg Ser Ser Gln Asn Ile Ile His Ser Asn Gly Asn Thr Tyr Leu Glu  
1 5 10 15

<210> 2

<211> 7

<212> PRT

<213> mus sp.

<400> 2

Lys Val Ser Asn Arg Phe Ser  
1 5

<210> 3

<211> 9

<212> PRT

<213> Mus sp.

<400> 3

Phe Gln Gly Ser His Val Pro Leu Thr  
1 5

<210> 4

<211> 7

<212> PRT

<213> Mus sp.

<400> 4

Thr Ser Gly Met Gly Val Ser  
1 5

<210> 5

<211> 16

<212> PRT

<213> Mus sp.

<400> 5

His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser Leu Lys Ser  
1 5 10 15

<210> 6

<211> 13

<212> PRT

<213> Mus sp.

<400> 6

Arg Pro Ile Thr Pro Val Leu Val Asp Ala Met Asp Tyr  
1 5 10

<210> 7

<211> 113

<212> PRT

<213> humanized antibody

<220>

<221> MISC\_FEATURE

<222> (3)..(3)

<223> Xaa=Val or Leu

<220>

<221> MISC\_FEATURE

<222> (7)..(7)

<223> Xaa=Ser or Thr

<220>

<221> MISC\_FEATURE

<222> (14)..(14)

<223> Xaa=Thr or Ser

<220>

<221> MISC\_FEATURE

<222> (17)..(17)

<223> Xaa=Gln, Asp, or Asn

<220>

<221> MISC\_FEATURE

<222> (30)..(30)

<223> Xaa=Ile or Val

<220>

<221> MISC\_FEATURE

<222> (50)..(50)

<223> Xaa=Arg or Lys

<220>

<221> MISC\_FEATURE

<222> (88)..(88)

<223> Xaa=Val or Leu

<220>

<221> MISC\_FEATURE

<222> (105)..(105)

<223> Xaa=Gly or Ala

<220>

<221> MISC\_FEATURE

<222> (109)..(109)

<223> Xaa=Val or Leu

<400> 7

Asp Val Xaa Met Thr Gln Xaa Pro Leu Ser Leu Pro Val Xaa Leu Gly  
1 5 10 15

Xaa Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Asn Ile Xaa His Ser  
20 25 30

Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser  
35 40 45

Pro Xaa Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro  
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Xaa Gly Val Tyr Tyr Cys Phe Gln Gly  
85 90 95

Ser His Val Pro Leu Thr Phe Gly Xaa Gly Thr Lys Xaa Glu Ile Lys  
100 105 110

Arg

<210> 8

<211> 123

<212> PRT

<213> humanized antibody

<220>

<221> MISC\_FEATURE

<222> (1)..(1)

<223> Xaa=Gln or Glu

<220>

<221> MISC\_FEATURE

<222> (2)..(2)

<223> Xaa=Val or Ala

<220>

<221> MISC\_FEATURE

<222> (64)..(64)

<223> Xaa=Ser or Thr

<220>

<221> MISC\_FEATURE

<222> (77)..(77)

<223> Xaa=Lys or Arg

<220>

<221> MISC\_FEATURE

<222> (78)..(78)

<223> Xaa=Ser or Thr

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (83)..(83)

&lt;223&gt; Xaa=Thr or Ser

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (84)..(84)

&lt;223&gt; Xaa=Met, Ile or Leu

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (86)..(86)

&lt;223&gt; Xaa=Asn, Ser, Thr

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (87)..(87)

&lt;223&gt; Xaa=Met, Val, or Leu

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (118)..(118)

&lt;223&gt; Xaa=Leu or Ser

&lt;400&gt; 8

Xaa Xaa Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu  
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser  
20 25 30

Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu  
35 40 45

Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Xaa  
50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Xaa Xaa Gln Val  
65 70 75 80

Val Leu Xaa Xaa Thr Xaa Xaa Asp Pro Val Asp Thr Ala Thr Tyr Tyr  
85 90 95

Cys Val Arg Arg Pro Ile Thr Pro Val Leu Val Asp Ala Met Asp Tyr  
100 105 110

Trp Gly Gln Gly Thr Xaa Val Thr Val Ser Ser  
115 120

<210> 9

<211> 113

<212> PRT

<213> humanized antibody

<400> 9

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly  
1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Asn Ile Ile His Ser  
20 25 30

Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser  
35 40 45

Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro  
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Phe Gln Gly  
85 90 95

Ser His Val Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys  
100 105 110

Arg

&lt;210&gt; 10

&lt;211&gt; 123

&lt;212&gt; PRT

&lt;213&gt; humanized antibody

&lt;400&gt; 10

Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu  
 1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser  
 20 25 30

Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu  
 35 40 45

Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser  
 50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Ser Gln Val  
 65 70 75 80

Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr  
 85 90 95

Cys Val Arg Arg Pro Ile Thr Pro Val Leu Val Asp Ala Met Asp Tyr  
 100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser  
 115 120

&lt;210&gt; 11

&lt;211&gt; 219

&lt;212&gt; PRT

&lt;213&gt; humanized antibody

&lt;400&gt; 11

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly  
 1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Asn Ile Ile His Ser  
 20 25 30



9/19

Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser  
 35 40 45

Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro  
 50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
 65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Phe Gln Gly  
 85 90 95

Ser His Val Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys  
 100 105 110

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu  
 115 120 125

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe  
 130 135 140

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln  
 145 150 155 160

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser  
 165 170 175

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu  
 180 185 190

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser  
 195 200 205

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
 210 215

<210> 12

<211> 453

<212> PRT

<213> humanized antibody

<400> 12

Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu  
 1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser  
 20 25 30  
 Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu  
 35 40 45  
 Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser  
 50 55 60  
 Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Ser Gln Val  
 65 70 75 80  
 Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr  
 85 90 95  
 Cys Val Arg Arg Pro Ile Thr Pro Val Leu Val Asp Ala Met Asp Tyr  
 100 105 110  
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly  
 115 120 125  
 Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly  
 130 135 140  
 Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val  
 145 150 155 160  
 Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe  
 165 170 175  
 Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val  
 180 185 190  
 Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val  
 195 200 205  
 Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys  
 210 215 220  
 Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu  
 225 230 235 240  
 Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr  
 245 250 255  
 Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val  
 260 265 270

11/19

Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val  
 275 280 285

Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser  
 290 295 300

Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu  
 305 310 315 320

Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala  
 325 330 335

Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro  
 340 345 350

Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln  
 355 360 365

Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala  
 370 375 380

Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr  
 385 390 395 400

Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu  
 405 410 415

Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser  
 420 425 430

Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser  
 435 440 445

Leu Ser Pro Gly Lys  
 450

<210> 13

<211> 46

<212> DNA

<213> DNA primer

<400> 13  
 tatagagctc aagcttggat ggtgggaaga tggatacagt tgggtgc

46

<210> 14

<211> 50

<212> DNA

<213> DNA primer

<400> 14

tatagagctc aagcttccag tggatagach gatggggstg tygttttggc 50

<210> 15

<211> 1953

<212> DNA

<213> humanized antibody

<400> 15

acgcgtccac catgaagttg cctgttaggc tgtttggtact gatgttcttg attcctgctt 60  
ccagcagtga tgttgtgatg acccaatctc cactctccct gcctgtcact cttggacagc 120  
cagcctccat ctcttcgaga tctagtcaga acattataca tagtaatgga aacacctatt 180  
tagaatggta cctgcagaaa ccaggccagt ctccaaggct cctgatctac aaagtttcca 240  
accgattttc tgggggtccca gacaggttca gtggcagtgg atcagggaca gatttcacac 300  
tcaagatcag cagagtggag gctgaggatg tgggagttta ttactgcttt caaggttcac 360  
atgttccgct cactttcggc ggagggacca aggtggaaat aaaacgtaag tgcactttcc 420  
taatctagaa attctaaact ctgaggggggt cggatgacgt ggccattctt tgcctaaagc 480  
attgagttta ctgcaaggct agaaaagcat gcaaagccct cagaatggct gcaaagagct 540  
ccaacaaaac aatttagaac ttatttaagg aataggggga agctaggaag aaactcaaaa 600  
catcaagatt ttaaatacgc ttcttgggtc ctttgctata attatctggg ataagcatgc 660  
tgttttctgt ctgtccctaa catgccctgt gattatccgc aaacaacaca cccaagggca 720  
gaactttgtt acttaaacac catcctgttt gcttctttcc tcaggaactg tggctgcacc 780  
atctgtcttc atcttcccgc catctgatga gcagttgaaa tctggaactg cctctgttgt 840  
gtgcctgctg aataacttct atcccagaga ggccaaagta cagtggaagg tggataacgc 900  
cctccaatcg ggtaactccc aggagagtgt cacagagcag gacagcaagg acagcaccta 960  
cagcctcagc agcaccctga cgctgagcaa agcagactac gagaaacaca aagtctacgc 1020  
ctgcgaagtc acccatcagg gcctgagctc gcccgtcaca aagagcttca acaggggaga 1080  
gtgttagagg gagaagtgcc cccacctgct cctcagttcc agcctgacct cctcccatcc 1140  
tttggcctct gacccttttt ccacagggga cctacccta ttgcggtcct ccagctcatc 1200  
tttcacctca cccccctcct cctccttggc ttttaattatg ctaatgttgg aggagaatga 1260

ataaataaag tgaatctttg cacctgtggg ttctctcttt cctcatttaa taattattat 1320  
ctgttgTTTT accaactact caatttctct tataagggac taaatatgta gtcacctaata 1380  
ggcgcataac cttttataaa aatcatcctt cattctatTT taccctatca tcctctgcaa 1440  
gacagtcctc cctcaaacc acaagccttc tgcctcaca gtcccctggg ccatggtagg 1500  
agagacttgc ttcttgTTT tcccctcctc agcaagccct catagtcctt ttttaagggtg 1560  
acaggcttta cagtcataata tcctttgatt caattccctg agaatcaacc aaagcaaatt 1620  
tttcaaaaga agaaacctgc tataaagaga atcattcatt gcaacatgat ataaaataac 1680  
aacacaataa aagcaattaa ataaacaaac aatagggaaa tgtttaagtt catcatggta 1740  
cttagactta atggaatgtc atgccttatt tacatTTTT aacagggtact gagggactcc 1800  
tgtctgccaa gggccgtatt gagtactttc cacaacctaa tttaatccac actatactgt 1860  
gagattaaaa acattcatta aaatgttgca aaggttctat aaagctgaga gacaaatata 1920  
ttctataact cagcaatccc acttctagga tcc 1953

<210> 16

<211> 3256

<212> DNA

<213> humanized antibody

<400> 16

acgcgtccac catggacagg cttacttcct cattcctgct gctgattgtc cctgcatatg 60  
tcctgtccca ggttactctg aaagagtctg gccctgtact agtgaagccc accgagaccc 120  
tcactctgac ttgtactttc tctgggtttt cactgagcac ttctggtatg ggagtgaact 180  
ggattcgtca gcctccagga aaggctctgg agtggtctgc acacatttac tgggatgatg 240  
acaagcgcta taaccatcc ctgaagagcc ggctcacaat ctccaaggat acctccaaaa 300  
gccaggtagt cctcacgatg accaatatgg accctgtaga tactgccaca tactactgtg 360  
ttcgaaggcc cattactccg gtactagtcg atgctatgga ctactggggc caaggaacct 420  
tggtcacctg ctctcaggt gagtcctcac aacctctaga gctttctggg gcaggccagg 480  
cctgaccttg gctttggggc agggaggggg ctaagggtgag gcagggtggc ccagccagg 540  
gcacacccaa tgcccatgag cccagacact ggacgtgaa cctcgcggac agttaagaac 600  
ccaggggcct ctgcgccctg ggcccagctc tgtcccacac cgcggtcaca tggcaccacc 660  
ttctttgag cctccaccaa gggcccatcg gtcttcccc tggcaccctc ctccaagagc 720  
acctctgggg gcacagcggc cctgggctgc ctggtcaagg actacttccc cgaaccgggtg 780  
acggtgtcgt ggaactcagg cgccctgacc agcggcgtgc acaccttccc ggctgtccta 840

cagtcctcag gactctactc cctcagcagc gtggtgaccg tgccctccag cagcttgggc 900  
accagacctt acatctgcaa cgtgaatcac aagcccagca acaccaaggt ggacaagaaa 960  
gttggtgaga ggccagcaca gggagggagg gtgtctgctg gaagccaggc tcagcgtcc 1020  
tgccctggacg catcccggct atgcagcccc agtccagggc agcaaggcag gccccgtctg 1080  
cctcttcacc cggaggcctc tgcccgtccc actcatgctc agggagaggg tcttctggct 1140  
ttttccccag gctctgggca ggcacaggct aggtgccctt aaccaggcc ctgcacacaa 1200  
aggggcagggt gctgggtca gacctgcaa gagccatata cgggaggacc ctgcccctga 1260  
cctaagccca ccccaaaggc caaactctcc actccctcag ctcgacacc ttctctctc 1320  
ccagattcca gtaactccca atcttctctc tgcagagccc aaatcttggtg aaaaaactca 1380  
cacatgcca cgtgcccag gtaagccagc ccaggcctcg ccctccagct caaggcggga 1440  
cagggtgccct agagtagcct gcatccaggg acaggcccca gccgggtgct gacacgtcca 1500  
cctccatctc ttctcagca cctgaactcc tggggggacc gtcagtctc ctcttcccc 1560  
caaaaccaa ggacaccctc atgatctccc ggaccctga ggtcacatgc gtggtggtgg 1620  
acgtgagcca cgaagaccct gaggtcaagt tcaactggta cgtggacggc gtggaggtgc 1680  
ataatgcca gacaaagccg cgggaggagc agtacaacag cacgtaccgt gtggtcagcg 1740  
tcctcaccgt cctgcaccag gactggctga atggcaagga gtacaagtgc aaggctctca 1800  
acaaagccct ccagcccc atcgagaaaa ccatctccaa agccaaaggt gggaccctgt 1860  
gggtgcgagg gccacatgga cagaggccgg ctcgccccc cctctgccct gagagtgacc 1920  
gctgtacca cctctgtccc tacagggcag ccccgagaac cacaggtgta caccctgccc 1980  
ccatcccggg atgagctgac caagaaccag gtcagcctga cctgcctggt caaaggcttc 2040  
tatcccagcg acatcgccgt ggagtgggag agcaatgggc agccggagaa caactacaag 2100  
accacgcctc ccgtgtgga ctccgacggc tcttcttcc tctacagca gctcaccgtg 2160  
gacaagagca ggtggcagca ggggaacgtc ttctcatgct ccgtgatgca tgaggctctg 2220  
cacaaccact acacgcagaa gagcctctcc ctgtctccgg gtaaagtagt gcgacggccg 2280  
gcaagcccc gctccccggg ctctcgcggt cgcacgagga tgcttggcac gtacccctg 2340  
tacatacttc cggggcggc agcatggaaa taaagcacc agcgctgccc tggggccctg 2400  
cgagactgtg atggttcttt ccacgggtca ggccgagtct gaggcctgag tggcatgagg 2460  
gaggcagagc ggggtccact gtccccacac tggcccaggc tgtgcagggtg tgccctgggcc 2520  
gcctaggggtg gggctcagcc aggggctgcc ctcggcaggg tgggggattt gccagcgtgg 2580  
ccctccctcc agcagcacct gccctgggct gggccacggg aagccctagg agccctggg 2640  
gacagacaca cagcccctgc ctctgtagga gactgtcctg ttctgtgagc gccctgtcct 2700  
ccgacctcca tgccactcg ggggcatgcc tagtccatgt gcgtagggac aggcctccc 2760

```

tcacccatct acccccacgg cactaacccc tggctgccct gccagcctc gcacccgcat 2820
ggggacacaa ccgactccgg ggacatgcac tctcggggccc tgtggaggga ctggtgcaga 2880
tgcccacaca cacactcagc ccagacccgt tcaacaaacc ccgcactgag gttggccggc 2940
cacacggcca ccacacacac acgtgcacgc ctcaacacg gagcctcacc cgggcgaact 3000
gcacagcacc cagaccagag caaggctctc gcacacgtga acactcctcg gacacaggcc 3060
cccacgagcc ccacgcggca cctcaaggcc cagagcctc tcggcagctt ctccacatgc 3120
tgacctgctc agacaaaccc agccctcctc tcacaagggg gccctgcag ccgccacaca 3180
cacacagggg atcacacacc acgtcacgtc cctggccctg gccacttcc cagtgccgcc 3240
cttcctgca ggatcc 3256

```

&lt;210&gt; 17

&lt;211&gt; 714

&lt;212&gt; DNA

&lt;213&gt; humanized antibody

```

<400> 17
atgaagttgc ctgttaggct gttggtactg atgttctgga ttcctgcttc cagcagtgat 60
gttgtgatga cccaatctcc actctccctg cctgtcactc ttggacagcc agcctccatc 120
tcttgagat ctagtcagaa cattatacat agtaatggaa acacctattt agaatggtac 180
ctgcagaaac caggccagtc tccaaggctc ctgatctaca aagtttccaa ccgattttct 240
ggggtcccag acaggttcag tggcagtggg tcaggagacag atttcacact caagatcagc 300
agagtggagg ctgaggatgt gggagtttat tactgctttc aaggttcaca tgttccgctc 360
actttcggcg gagggaccaa ggtggaaata aaacgaactg tggctgcacc atctgtcttc 420
atcttcccg cactctgatga gcagttgaaa tctggaactg cctctgttgt gtgcctgctg 480
aataacttct atcccagaga ggccaaagta cagtggaagg tggataacgc cctccaatcg 540
ggtaactccc aggagagtgt cacagagcag gacagcaagg acagcaccta cagcctcagc 600
agcaccctga cgctgagcaa agcagactac gagaaacaca aagtctacgc ctgcgaagtc 660
acccatcagg gcctgagctc gcccgtcaca aagagcttca acaggggaga gtgt 714

```

&lt;210&gt; 18

&lt;211&gt; 1416

&lt;212&gt; DNA

&lt;213&gt; humanized antibody

<400> 18  
atggacaggc ttacttcctc attcctgctg ctgattgtcc ctgcatatgt cctgtcccag 60  
gttactctga aagagtctgg ccctgtacta gtgaagccca ccgagaccct cactctgact 120  
tgtactttct ctgggttttc actgagcact tctggtatgg gagtgagctg gattcgctag 180  
cctccaggaa aggctctgga gtggctggca cacatttact gggatgatga caagcgctat 240  
aaccatccc tgaagagccg gctcacaatc tccaaggata cctccaaaag ccaggtagtc 300  
ctcacgatga ccaatatgga ccctgtagat actgccacat actactgtgt tcgaaggccc 360  
attactccgg tactagtcca tgctatggac tactggggcc aaggaaccct ggtcaccgtc 420  
tcctcagcct ccaccaaggg cccatcggtc tccccctgg caccctcctc caagagcacc 480  
tctgggggca cagcggccct gggctgcctg gtcaaggact acttccccga accggtgacg 540  
gtgtcgtgga actcaggcgc cctgaccagc ggcgtgcaca ccttccccgc tgtcctacag 600  
tcctcaggac tctactccct cagcagcgtg gtgaccgtgc cctccagcag cttgggcacc 660  
cagacctaca tctgcaacgt gaatcacaag cccagcaaca ccaagggtgga caagaaagtt 720  
gagcccaaat cttgtgacaa aactcacaca tgcccaccgt gcccagcacc tgaactcctg 780  
gggggaccgt cagtcttctc cttcccccca aaaccaagg acaccctcat gatctcccgg 840  
acccctgagg tcacatgcgt ggtggtggac gtgagccacg aagaccctga ggtcaagttc 900  
aactggtacg tggacggcgt ggaggtgcat aatgccaaga caaagccgcg ggaggagcag 960  
tacaacagca cgtaccgtgt ggtcagcgtc ctcaccgtcc tgcaccagga ctggctgaat 1020  
ggcaaggagt acaagtgcaa ggtctccaac aaagccctcc cagcccccat cgagaaaacc 1080  
atctccaaag ccaaagggca gccccgagaa ccacaggtgt acaccctgcc cccatcccgg 1140  
gatgagctga ccaagaacca ggtcagcctg acctgcctgg tcaaaggctt ctatcccagc 1200  
gacatcgccg tggagtggga gagcaatggg cagccggaga acaactacaa gaccacgcct 1260  
cccgtgctgg actccgacgg ctcttcttct ctctacagca agctcaccgt ggacaagagc 1320  
aggtggcagc aggggaacgt cttctcatgc tccgtgatgc atgaggctct gcacaaccac 1380  
tacacgcaga agagcctctc cctgtctccg ggtaaa 1416

<210> 19

<211> 238

<212> PRT

<213> humanized antibody

<400> 19



Met Lys Leu Pro Val Arg Leu Leu Val Leu Met Phe Trp Ile Pro Ala  
 1 5 10 15  
 Ser Ser Ser Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val  
 20 25 30  
 Thr Leu Gly Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Asn Ile  
 35 40 45  
 Ile His Ser Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro  
 50 55 60  
 Gly Gln Ser Pro Arg Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser  
 65 70 75 80  
 Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr  
 85 90 95  
 Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys  
 100 105 110  
 Phe Gln Gly Ser His Val Pro Leu Thr Phe Gly Gly Gly Thr Lys Val  
 115 120 125  
 Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro  
 130 135 140  
 Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu  
 145 150 155 160  
 Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn  
 165 170 175  
 Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser  
 180 185 190  
 Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala  
 195 200 205  
 Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly  
 210 215 220  
 Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys  
 225 230 235

&lt;210&gt; 20

&lt;211&gt; 472

&lt;212&gt; PRT

&lt;213&gt; humanized antibody

&lt;400&gt; 20

Met Asp Arg Leu Thr Ser Ser Phe Leu Leu Leu Ile Val Pro Ala Tyr  
 1 5 10 15

Val Leu Ser Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys  
 20 25 30

Pro Thr Glu Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu  
 35 40 45

Ser Thr Ser Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys  
 50 55 60

Ala Leu Glu Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr  
 65 70 75 80

Asn Pro Ser Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys  
 85 90 95

Ser Gln Val Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala  
 100 105 110

Thr Tyr Tyr Cys Val Arg Arg Pro Ile Thr Pro Val Leu Val Asp Ala  
 115 120 125

Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser  
 130 135 140

Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr  
 145 150 155 160

Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro  
 165 170 175

Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val  
 180 185 190

His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser  
 195 200 205

Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile  
 210 215 220

Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val  
 225 230 235 240  
 Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala  
 245 250 255  
 Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro  
 260 265 270  
 Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val  
 275 280 285  
 Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val  
 290 295 300  
 Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln  
 305 310 315 320  
 Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln  
 325 330 335  
 Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala  
 340 345 350  
 Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro  
 355 360 365  
 Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr  
 370 375 380  
 Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser  
 385 390 395 400  
 Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr  
 405 410 415  
 Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr  
 420 425 430  
 Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe  
 435 440 445  
 Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys  
 450 455 460  
 Ser Leu Ser Leu Ser Pro Gly Lys  
 465 470